

Full Length Research Paper

Exploring carotenoid levels and characteristics in fish fillets from West African coastal fisheries

Bazyli Czeczuga^{1*}, Janusz Semeniuk², Ewa Czeczuga-Semeniuk¹, Adrianna Semeniuk¹ and Bernard Klyszejko³

¹Department of General Biology, Medical University, Mickiewicza 2C, 15-222 Białystok, Poland.

²Department of Pediatrics, Pediatric Gastroenterology and Allergology, Medical University, Waszyngtona 17, 15-274 Białystok, Poland.

³Department of Fish Physiology, Agricultural University, K. Krolewicz 4, 71-550 Szczecin, Poland.

Accepted 8 October, 2023

Using column (CC), thin- layer (TLC) and high- performance liquid chromatography (HPLC), carotenoid content was examined in the fillets (muscles with skin) of 16 fish species from the fisheries of West African Coast. 15 carotenoids, including 6 ketocarotenoids (4'- hydroxyechinenone, canthaxanthin, phoenicopterone, phoenicoxanthin, astaxanthin, 7,8- didehydroastaxanthin) were found, with a predominance of 3,4- dihydro- α - carotene, zeaxanthin, phoenicopterone, canthaxanthin, phoenicoxanthin and astaxanthin. Total carotenoid content in the examined material ranged from 0.086 (*Merluccius merluccius*) to 0.352 $\mu\text{g g}^{-1}$ wet mass (*Macrurus aequalis*). Also, the increase of different environmental factors- especially food, parasites and organochlorine pollution on carotenoid content in fishes and their role in human's life was discussed.

Key words: Carotenoids, fish, fillets, fisheries, West Africa.

INTRODUCTION

Carotenoids are wide spread and important pigments in nature. They occur in all families of flora and fauna. Only bacteria, fungi and plants are able to synthesize them *de novo*; animals have to obtain them from food. Fish meat owes its nutrient value not only to protein and fats, but also to other biologically active substances- including carotenoids, which plays an important role in humans' life. Carotenoids serve as a source of vitamin A, antioxidant and pro- oxidant (Yeum et al., 2009), play a cancer- protective role (Rock, 2009) and increase the immune response in mammals (Chew and Park, 2009). Carotenoids may also be significant in the coronary heart disease (Johnson and Krynski, 2009). Therefore, many authors have been carrying out extensive studies on the

presence and biological function of carotenoids in plants, animals and in humans. Thus, the knowledge about the carotenoid content in meat or respective fish species in fisheries seems to be of a great importance.

We have already investigated some of the species from the Baltic Sea (Czeczuga and Klyszejko, 1996), Black Sea (Czeczuga, 1973), fishing areas of the Antarctic (Czeczuga, 1978 a,b; 1982; Czeczuga and Klyszejko, 1978, 1986) and from the fisheries of New Zealand (Czeczuga et al., 2000). Tsukuda and Amono (1966), Matsuno et al. (1974, 1979), Matsuno and Katsuzama (1976) and Miki et al. (1982) have investigated the carotenoids in species from the waters washing the Japanese islands. Later, the analyses of carotenoids found in some species from the ocean near the coasts of California have been made (Crozier, 1967). Tanaka et al. (1978) have investigated the carotenoids in marine yellow fish from tropical regions. Carotenoids content in particular species

*Corresponding author. E-mail: bazylio@poczta.onet.pl.

of Pacific salmon have also been investigated (Jarzombek, 1970; Czczuga, 1979; Kitahara, 1984) and some species from an ocean ranching farm near Islands they were analysed by Czczuga et al. (2005).

Reviewed papers about carotenoids in fishes are generally dated prior to 2000 (Fox, 1957; Simpson et al., 1981); more recent works cover the carotenoids in general (Bjerkeng, 2008).

The present paper discusses the results of the analysis of carotenoid content in fishes from some fisheries of West African Coast.

MATERIAL AND METHODS

The study population included 16 fish species (Table 2) from the fisheries of the West African Coast (17°05'–22°35'S; 0.11°20'–0.13°30'E southern Africa and 22°0'–17°10'E northern Africa), caught in July and August.

50 g fillets (muscles with skin) have been used in the investigations. We analysed the material from three specimens. After three weeks of storage, the refrigerated material (–4°C) which has been used for the analysis, was sent by air mail to the department laboratory where they were analysed a week later. The carotenoid pigments were isolated using column (CC), thin-layer (TLC) and high-performance liquid chromatography (HPLC).

Prior to chromatography, the material was homogenized and hydrolyzed for 24 h in a 10% solution of nitrogen, at room temperature. The extract was subsequently placed onto an Al₂O₃-filled Quickfit Co. column. The individual fractions were eluted using various solvent systems (Czczuga, 1988). The eluent was evaporated and the residue was dissolved in appropriate solvent to draw the maximum of absorption. This was necessary, among other reasons, to identify particular carotenoids. In addition to column chromatography, the acetone extract has been divided into fractions with thin-layer chromatography. Silicon-gel-covered glass plates (Merck Co.) and various solvent systems were used. The R_f values were established according to commonly accepted criteria (Schiedt and Liaen-Jensen, 1995).

Pigments were also determined by ion-pairing in reverse-phase HPLC. The HPLC equipment consisted of a Shimadzu SCL-6B gradient programmer and a Reodyne 7125 injector equipped with a 20 µl loop. Detection was achieved in a Shimadzu SPD-6AV UV-VIS spectrophotometric detector set at 440 nm and a Shimadzu RF-535 fluorescence detector.

Carotenoids were identified through comparison with the standards from: a) the behaviour in CC; b) their UV-UIS spectra; c) their partition between n-hexane and 95% ethanol; d) their R_f-values in TLC; e) the presence of the allylic OH group determined by the acid CHCl₃ test; f) the epoxide test; g) the mass spectrum (Vetter et al., 1971).

Carotenoid pigment standards were purchased from the Hoffman-La Roche Company, Switzerland, the International Agency for the ¹⁴C Determinations, Denmark, and Sigma Chemical Company, USA.

Carotenoids recorded in investigated fishes belong to three groups (hydrocarbon, hydroxycarotenoids and ketocarotenoids) which chemical, structural and semisystematical characteristics have been placed in monographs of Foppen (1971), Isler (1971) and Straub (1987). We have also used those monographs in our investigations. Quantitative analyses were performed with UV-UIS spectroscopy according to Davies methods (Czczuga, 1988). The structure of carotenoids was described by Straub (1987). Chromatography methods (CC, TLC, HPLC) were described in detail by Bernhard (1995), Schiedt (1995) and Pfander and Riesen (1995) respectively. The results were evaluated with Scheffe Test (Winer, 1997).

RESULTS AND DISCUSSION

Fifteen (15) carotenoids were found in the examined material (Table 1, Figure 1). Most of them were common in fish, but some, such as 3,4-didehydro-α-carotene, 4-hydroxy-α-carotene, phoenicopterone and 7,8-didehydroastaxanthin were rather rare. In most of the species, lutein, zeaxanthin, phoenicopterone and astaxanthin were found (Table 2). In the investigated material, 3,4-didehydro-α-carotene, zeaxanthin, phoenicopterone, phoenicoxanthin, canthaxanthin and astaxanthin were the predominant group. The total carotenoid content in the investigated fish species ranged from 0.086 (*Merluccius merluccius*) to 0.424 µg g^{–1} wet mass (*Pterothrissus belloci*). Mean value for all investigated species averaged 0.200 µg g^{–1} wet mass.

The total carotenoid content in the fillets of the species from African fishing areas in comparison with species from other fisheries were considered (mean 0.200 µg g^{–1} wet mass). The mean value for nine species from the region of the Falkland Islands was 0.08 (Czczuga and Klyszejko, 1978) and for 10 species from the Szczecin Lagoon of Baltic Sea was 0.230 µg g^{–1} wet mass of fillets (Czczuga and Klyszejko, 1996). Similar values were observed in fillets of fishes from Black Sea fisheries (Czczuga, 1973), New Zealand fishing areas (Czczuga et al., 2000) and from the waters washing the Japanese Islands (Tsukuda and Amono, 1966; Matsuno et al., 1974; Matsuno and Katsuyama, 1976; Miki et al., 1982).

In fillets from Antarctic fish species, especially in representatives of Nototheniidae and Chaenichthyidae (white blooded fish) families, total carotenoid content was low (Czczuga, 1978a, b; 1982). In Atlantic (Torrisen et al., 1989; Czczuga et al., 2005) and Pacific salmonids (Jarzombek, 1970; Crozier, 1970; Kitahara, 1984), the biggest amounts of carotenoids are retained in the fillets.

Apart from carotenoids commonly occurring in fishes, some rare types have also been found. One of them is 3,4-didehydro-α-carotene. It was first isolated from the body of sheatfish *Silurus glanis* (Czczuga, 1977). Although it has been found in material from six examined species, only in *Merluccius merluccius* was it a predominant carotenoid. 4-hydroxy-α-carotene- also called β,ε-carotene-4-ol, was first observed in plant material by Zechmeister in 1958 (Isler, 1971). It also occurred in sheatfish (Czczuga, 1977a). We found it only in 2 of the 16 investigated species; in *Belone belone* and *Centrolophus niger*. Phoenicopterone and 4-keto-α-carotene derivative is frequently called 4-keto-α-carotene. It was observed in nine species and in *Belone belone* and *Dentex macropthalmus* specimens as predominant carotenoid. Phoenicopterone was found for the first time in certain green algae as extra-plastidic pigment (Goodwin, 1980). 7,8-didehydroastaxanthin, astaxanthins' derivative, was being frequently called asterinic acid in previous studies and has been noted only in *Belone belone* and *Clupea pilchardus* individuals. This acetylenic Zeaxanthin occurs in fillets of investigated 13

Table 1. Carotenoid list from investigated material

Carotenoid	Summary formula	Structure (Figure 1)	Semisystematic name
α - Carotene	C ₄₀ H ₅₆	A-R-B	β,ϵ - Carotene
β - Carotene	C ₄₀ H ₅₆	B-R-B	β,β - Carotene
ϵ - Carotene	C ₄₀ H ₅₆	A-R-A	ϵ,ϵ - Carotene
3,4- Didehydro- α - carotene	C ₄₀ H ₅₄	A-R-C	3,4- Didehydro- β,ϵ - carotene
β - Cryptoxanthin	C ₄₀ H ₅₆ O	B-R-D	β,β - Carotene- 3- ol
4- Hydroxy- α - carotene	C ₄₀ H ₅₆ O	A-R-E	β,ϵ - Carotene- 4- ol
Lutein	C ₄₀ H ₅₆ O ₂	F-R-D	β,ϵ - Carotene- 3,3'- diol
Tunaxanthin	C ₄₀ H ₅₆ O ₂	F-R-F	ϵ,ϵ - Carotene- 3,3'- diol
Zeaxanthin	C ₄₀ H ₅₆ O ₂	D-R-D	β,β - Carotene- 3,3'- diol
Phoenicopterone	C ₄₀ H ₅₄ O	A-R-G	β,ϵ - Carotene- 4- ol
4'- Hydroxyechinenone	C ₄₀ H ₅₄ O ₂	E-R-G	4'- Hydroxy- β,β - caroten- 4-one
Canthaxanthin	C ₄₀ H ₅₂ O ₂	G-R-G	β,β - Carotene- 4,4'- dione
Phoenicoxanthin	C ₄₀ H ₅₂ O ₃	G-R-H	3- Hydroxy- β,β - carotene- 4,4'-dione
Astaxanthin	C ₄₀ H ₅₂ O ₄	H-R-H	3,3'- Dihydroxy- β,β - carotene- 4,4'-dione
7,8- Didehydroastaxanthin	C ₄₀ H ₅₀ O ₄	I-R ₁ -H	3,3'- Dihydroxy- 7,8- dihydro- β,β - carotene- 4,4'-dione




Figure 1. Structural features of carotenoids from the analysed materials. A- I, end group designation of carotenoids; R, R₁ polyene chain.

Table 2. Carotenoid content in the investigated material.

Specie	Carotenoid (Table 1)	Content of particular carotenoids (%)	Total content ($\mu\text{g g}^{-1}$ wet mass) $\pm$ SD
<i>Acanthias acanthias</i> L.	2,3,7,8,9,14	9 (33.6), 2(21.3), 14 (19.2), 7 (9.5), 8(9.5), 3 (7.5)	0.089
<i>Belone belone</i> (Brünn.)	6,7,9,10,15	10 (21.9), 9 (21.5), 15 (20.9), 6 (18.4), 7 (17.3)	0.143
<i>Centrolophus niger</i> (Gmelin)	5,6,7,11,12,14	12 (23.1), 11 (22.8), 14 (20.9), 5 (15.7), 6 (10.2), 7 (7.3)	0.121
<i>Clupea pilchardus</i> Walb.	7,9,10,14,15	14 (47.6), 10 (28.5), 15 (8.2), 9 (8.1), 7 (7.6)	0.259
<i>Dentex macroptalmus</i> (Bloch.)	7,8,10,14	10 (43.0), 7 (39.2), 14 (10.1), 8 (7.7)	0.153
<i>Engraulis encrasicolus</i> L.	2,3,7,9,10,14	14 (33.2), 10 (30.8), 2 (20.5), 3 (8.6), 7 (4.2), 9 (2.7)	0.258
<i>Genypterus capensis</i> (Smith.)	1,2,4,5,7,9,14	9 (22.6), 14 (21.8), 5 (18.4), 2 (15.0), 1 (12.4), 4 (6.8), 7 (3.0)	0.231
<i>Macrurus aequalis</i> (Ginter)	1,3,4,5,9,10,12	12 (19.4), 5 (18.3), 9 (16.5), 10 (14.9), 1 (12.8), 3 (11.6), 4 (6.5)	0.352
<i>Merluccius merluccius</i> L.	4,5,7,8,9,10,14	4 (26.7), 7 (24.8), 8 (21.4), 5 (18.8), 9 (4.4), 10 (2.7), 14 (1.2)	0.086
<i>Myliobatis aquila</i> L.	3,7,9,13,14	13 (36.0), 7 (28.2), 14 (25.4), 9 (7.6), 3 (2.8)	0.202
<i>Pterothrissus belloci</i> (Cadenat)	3,4,5,9,10,12	12 (34.4), 5 (20.6), 10 (19.2), 9 (18.6), 3 (5.4), 4 (1.8)	0.424
<i>Sarda sarda</i> (Bloch)	2,5,7,10,14	14 (44.5), 10 (30.2), 2 (18.4), 5 (6.2), 7 (0.7)	0.305
<i>Sebastes dactyloptera</i> (Dela Roche)	2,4,5,7,9,12,14	9 (28.6), 2 (26.4), 5 (18.8), 7 (12.2), 4 (10.3), 14 (2.5), 12 (1.2)	0.265
<i>Trachurus trachurus</i> L.	1,2,3,5,9,14	14 (22.8), 2 (21.4), 1 (20.8), 3 (18.4), 5 (14.2), 9 (2.4)	0.108
<i>Trigla lyra</i> L.	4,5,9,10,12,14	12 (24.2), 5 (22.2), 10 (21.4), 14 (20.4), 4 (8.2), 9 (3.6)	0.090
<i>Zeus faber</i> L.	2,7,8,9,14	14 (25.4), 2 (24.8), 7 (21.3), 9 (17.6), 8 (10.9)	0.111

species, while lutein was found in 12 of them. Zeaxanthin appears predominant in *Acanthias acanthias*, *Genypterus capensis*, and *Sebastes dactyloptera*.

These are the carotenoids belonging to the yellow group and are the principle carotenoids of the lens and the macula of humans and primate eye (Schalch et al., 2009). The role of this dihydroxy compounds in risk reduction of macular degeneration and cataract of the eye are important. From all of the environmental factors such ones as food (Simpson et al., 1981; Latscha, 1990) increases the carotenoid content in fish.

The studies included 17 species of invertebrates (food of fish) belonging to Porifera, Coelenterata, Annelida, Crustacea, Mollusca and Echinodermata which were caught along the coast of West Africa (Czeczuga and Klyszejko, 1977). The investigations showed that they have been the

richest in carotenoids. 27 particular carotenoids have been identified. Total carotenoid content in this material varied from 0.039 to 1.529 $\mu\text{g g}^{-1}$ fresh weight. The lowest amount of carotenoids was found in the representatives of Coelenterata and the highest in the representatives of Crustacea. In natural conditions, the total carotenoid content and the amount of keto- and dihydroxycarotenoids in the respective parts of the fishes' body depends not only on the type of food it consume but also on the parasites and the organochlorine pollutants. The effect of ecto- and edoparasites on the carotenoid content in fish in natural infection was also described (Czeczuga et al., 2009). This study was conducted using pairs of the host-parasite with ectoparasites and endoparasites. In all examined pairs of host and parasite, the parasitic organism has always had the higher total carotenoid content in comparison

to the host. Carotenoid content in healthy and infected fishes ranged from 3.1 (*Mollienisia latipinna*) to 51.9 (intestines of *Tinca tinca*). Obtained data indicates that some carotenoids are selectively accumulated by respective parasites.

Water polluted with organochlorine substances causes the so called M74 syndrome in fish, especially in salmon (Vuorinen et al., 1997). Afflicted are the females, especially the eggs which are pale yellow as they have a low red carotenoid content (especially low astaxanthin level) (Pettersson and Liguell, 1999; Czeczuga et al., 2002, 2005). Most of the larvae which develop from such eggs die when they begin the active feeding process. M74 syndrome occurs not only in Baltic salmon, but has also been noted in the sea trout *Salmo trutta* m. *trutta* (Czeczuga et al., 2005). The effect of chemical water pollution on the larval form of other fishes is well known in

other latitudes. The high mortality of early life- stage salmonids including Pacific salmon from some of the Great Lakes of North America has been reported under the name of Early Mortality Syndrome (EMS) (McDonald, 1995). Clinical symptoms were similar to those noted in M74 in salmon from the Baltic Sea or in EMS in other salmonid species from the North American Great Lakes; were observed in Atlantic salmon specimens with Cayuga syndrome in the New York Finger Lakes (Fisher et al., 1996). All three of those disorders responsible for mortality in early life- stage salmonids have been characterized (in females and eggs) through the low level of astaxanthin and thiamine (Fitzsimons et al., 1999).

Fish, like other animals, do not synthesize carotenoids *de novo*, which only plants, bacteria and fungi are able to synthesize. Those substances get to animals organisms only through food and some of them can be converted into other carotenoids via oxidation or reduction and into vitamin A. Carotenoids catered with food are being released in the intestines during digestion and are absorbed along the alimentary tract in fish (March et al., 1990). As revealed by studies on fish, the free vitamin A is formed not only from carotenoids which have three beta end groups (β - ring) from the β - carotene type (Simpson et al., 1981, Latscha, 1990), but also from a number of xanthophylls, including those from dihydroxy compounds- such as lutein, zeaxanthin or tunaxanthin (Katsuyama and Matsuno, 1987) as well as astaxanthin and canthaxanthin belonging to ketocarotenoids compounds (Guillou et al., 1989). Both, dihydroxy compounds and ketocarotenoids are quite common in fish species from the fisheries of Africa.

The shift of carotenoids in the pre- and postreproductive period occurs in fishes. At first, the shift has been reported only from the salmonids in the Pacific (Crozier, 1970; Kitahara, 1983). Further studies have showed, that this phenomenon is also typical to salmonid species from the *Salmo* genus (Czeczuga and Chelkowski, 1984) and freshwater fish species (Czeczuga and Czeczuga-Semeniuk, 2002). Differences are noted in the shift between sexes. Both, in males and females, the liver is the main reservoir of carotenoids, and on the second range, the intestines. In the females, in prespawning season, carotenoids mostly shift to the gonads and in males mainly to the skin and fins giving the mating colouration. A similar pattern occurs in amphibians (Czeczuga et al., 2006).

Conclusions

In the fillets of 16 fish species from the fisheries of West African coast, 15 carotenoids were found. Total carotenoids content ranged from 0.086 to 0.424 with mean value of $0.200 \mu\text{g g}^{-1}$ wet mass. Seven out of 15 carotenoids such as α - carotene, β - carotene, β - cryptoxanthin, lutein, zeaxanthin, canthaxanthin and astaxanthin play a significant role in the humans' health. They all have been

postulated to increase an immune (Chew and Park, 2009) and cancer protective role (Rock, 2009) in mammals. β - carotene and other carotenoids from this group behave as antioxidant factors at low oxygen pressure and as pro- oxidants at higher oxygen partial pressure (Yeum et al., 2009).

β - carotene and lutein have been postulated to have the antiproliferative effect on the cells (Palozza et al., 2009). Known are the effects of those carotenoids on the cell signalling and communication including cell cycle, apoptosis, cell differentiation and growth factors (Wang, 2009). Also, such derivatives of β - carotene as the β - apocarotenals are biologically active. Those carotenoid metabolites can offer protection against chronic diseases and certain cancers. The photoprotective effects of those carotenoids especially β - carotene towards skin damage induced by UVA and UVB have been reviewed (Goralczyk and Wertz, 2009). Concluding, α - carotene, β - carotene, β - cryptoxanthin, lutein and zeaxanthin may play a significant role in the coronary heart disease (Johnson and Krinsky, 2009). This all carotenoids are quite common in fish species from the fisheries of West Africa.

REFERENCES

- Bernhard K (1995). Chromatography: Part II. Column chromatography. In: Britton G et al. (Eds.). Carotenoids, vol. 1A: Isolation and analysis, Birkhäuser Verlag, Basel, Boston, Berlin, pp 117-130.
- Bjerkeng B (2008). Carotenoids in aquaculture: fish and crustaceans. In: Britton G et al. (Eds.). Carotenoids, vol. 4: Natural functions, Birkhäuser Verlag, Basel, Boston, Berlin, pp 237-254.
- Chew BP, Park JS (2009). The immune system. In: Britton G et al. (Eds.). Carotenoids, vol. 5: Nutrition and health, Birkhäuser Verlag, Basel, Boston, Berlin, pp 363-382.
- Crozier GF (1967). Carotenoids of seven species of *Sebastes*. Comp. Biochem. Physiol. 23:179-184.
- Crozier GF (1970). Tissue carotenoids in prespawning and spawning sockeye salmon (*Oncorhynchus nerka*). J. Fish. Res. Board Can. 27: 973-975.
- Czeczuga B (1973). Carotenoids and Vitamin A in some fishes from the coastal region of the Black Sea. Hydrobiol. 41: 113-125.
- Czeczuga B (1977). Carotenoids in fish *Silurus glanis* L. Pol. Arch. Hydrobiol. 24: 563-567.
- Czeczuga B (1978a). Carotenoids in fish. Representatives of family Nototheniidae from the Antarctic. Pol. Arch. Hydrobiol. 25: 705-709.
- Czeczuga B (1978b). Studies of carotenoids in white- blooded fish (Chaenichthyidae) from the Antarctic. Pol. Arch. Hydrobiol. 25: 711-715.
- Czeczuga B (1982). Carotenoids in uneven aged individuals of certain Antarctic species of fish. Acta Ichthyol. Piscat. 12: 107-116.
- Czeczuga B (1988). Carotenoids. In: Galun M (Ed.) CRC Handbook of Lichenology, vol. 3, CRC Press, Boca Raton, Florida, pp 25-34.
- Czeczuga B, Chelkowski Z (1984). Carotenoid content in adult individuals of sea trout *Salmo trutta* L. during spawning migration, spawning and post- spawning migration. Acta Ichthyol. Piscat. 14:187- 201.
- Czeczuga B, Czeczuga- Semeniuk E, Tryggvadottir SV (2005). Carotenoid content in various body parts of Atlantic salmon (*Salmo salar* L.) and Arctic charr (*Salvelinus alpinus* L.) specimens from an ocean ranching farm. Ocean. Hydrobiol. Stud. 34:31-42.
- Czeczuga B, Czeczuga- Semeniuk E (2002). Carotenoid content in *Anguilla anguilla* (L.) individuals undertaking spawning migration. Folia Biol. 48: 1-6.

- Czczuga B, Czczuga- Semenik E, Semenik A (2006). Changes in carotenoid content in the respective body parts of frogs specimens in the period of life activity. *Trands Comp. Biochem. Physiol.* 12:21-29.
- Czczuga B, Czczuga- Semenik E, Semenik A (2009). Effect of ecto- and endoparasites on the carotenoid content in fish in natural infection. *Trends Comp. Biochem. Physiol.* 14:55-64.
- Czczuga B, Klyszejko B (1976). Carotenoid content in invertebrates caught along the coast of West Africa. *Bull. Acad. Polon. Sci., Ser. Sci. biol.* 24: 719-723.
- Czczuga B, Klyszejko B (1978). Carotenoids in certain species of fish from Antarctic. *Hydrobiol.* 60:173-175.
- Czczuga B, Klyszejko B (1986). Carotenoids in fish from the Falklands region. *Acta Ichthyol. Piscat.* 16: 73-86.
- Czczuga B, Klyszejko B (1996). Abundance of carotenoids in fishes from the Szczecin Lagoon. *Zesz. Nauk. Akad. Rol. Szczecin* 171:3-10.
- Czczuga B, Klyszejko B, Czczuga- Semenik E (2000). The carotenoid content in certain fish species from the fisheries of New Zealand. *Bull. Sea Fisher. Inst.* 149:35-42.
- Czczuga B, Bartel R, Czczuga- Semenik E (2002). Carotenoid content in eggs of Atlantic salmon (*Salmo salar* L.) and brown trout (*Salmo trutta* L.) entering polish rivers for spawning or reared in fresh water. *Acta Ichthyol. Piscat.* 32:3-21.
- Czczuga B, Bartel R, Czczuga- Semenik E, Kosielinski P, Grochowski A (2005). The occurrence of M74 syndrome in sea trout (*Salmo trutta* m. *trutta* L.) individuals returning to spawn in the polish rivers of the Vistula catchment area. *Arch. Pol. Fish.* 13:39-49.
- Euler H von, Hellström H (1934). Über Asterinsäure, eine Carotenoisäure aus Seesternen. *Hoppe- Seyl. Z.* 223:89-97.
- Fisher JP, Bush B, Spitzbergen JM (1996). Contrasting pathologies associated with Cayuga syndrome and PCS induced mortality in early life stages of Atlantic salmon. *Swed. Environ. Protec. Agency Stockholm, Rep.* 4534:87-89.
- Fitzsimons JD, Brown SB, Honeyfield DC, Hnath JG (1999). A review of early mortality syndrome (EMS) in Great Lakes salmonids: relationship with thiamine deficiency. *Ambio* 28:9-15.
- Fontana JD, Czczuga B, Bonfim JMB, Chociai MB, Oliveira BH, Guimaraes MF, Baran M (1996). Bioproduction of carotenoids: the comparative use of raw sugarcane juice and depolymerised haggasse by *Phaffia rhodozyma*. *Biores. Technol.* 58:121-125.
- Foppen EH (1971). Tables for the identification of carotenoid pigments. *Chrom. Rev.* 14:133-298.
- Fox DL (1957). Carotenoids in fish. In: Brown ME (Ed.), *Physiology of Fishes*, Academic Press, New York, pp367-385.
- Goodwin TW (1980). *The Biochemistry of Carotenoids*, vol. 1 Plants, Chapman and Hall, London and New York, 377pp.
- Goralczyk R, Wertz K (2009). Skin photoprotection by carotenoids. In: Britton G et al. (Eds.), *Carotenoids*, vol. 5: Nutrition and health, Birkhäuser verlag, Basel, Boston, Berlin, pp 335-362.
- Guillou A, Choubert G, Storebakken T, De La Noüe J, Kaushik S (1989). Bioconversion pathway of astaxanthin into retinol, in mature rainbow trout (*Salmo gairdneri* Rich.). *Comp. Biochem. Physiol.* 94B:481-485.
- Isler O (1971). *Carotenoids*, Birkhäuser Verlag, Basel- Stuttgart, 932 pp.
- Jarzombek AA (1970). Carotenoids in salmonids and their role in reproduction. *Proc. WNIRO* 60:234-267.
- Johnson EJ, Krinsky NI (2009). Carotenoids and coronary heart disease. In: Britton G et al. (Eds.), *Carotenoids*, vol. 5: Nutrition and health, Birkhäuser verlag, Basel, Boston, Berlin, pp 287-300.
- Katsuyama M, Matsuno T (1987). Metabolism of three stereoisomers of astaxanthin in the fish, rainbow trout and tilapia. *Comp. Biochem. Physiol.* 86B: 1-5.
- Kitahara T (1983). Behaviour of carotenoids in the chum salmon (*Oncorhynchus keta*) during anadromous migration. *Comp. Biochem. Physiol.* 76B: 97-101.
- Kitahara T (1984). Carotenoids in Pacific salmon during the marine period. *Comp. biochem. Physiol.* 78B: 859-862.
- Latscha T (1990). Carotenoids- their nature and significance in animal feeds. *F. Hoffman- La Roche*, Basel, 342pp.
- March BE, Hajen WE, Deacon G, Mac Millan C, Walsh MG (1990). Intestinal absorption of astaxanthin, plasma astaxanthin concentration, body weight and metabolic rate as determinants of flesh pigmentation in salmonid fish. *Aquacult.* 90:313-322.
- Matsuno T, Katsuyama M (1976). Comparative biochemical studies of carotenoids in fishes 9. on the nineteen species of fishes in the division Percichthyes. *Bull. Jap. Soc. Sci. fish.* 42:645-649.
- Matsuno T, Matsutaka H, Katsuyama M, Nagata S (1979). Isolation of three stereoisomeric tunaxanthin A, B and C from tunaxanthin fraction in Japanese fishes. *Bull. Jap. Soc. Sci. Fish.* 45:1197-2009.
- Matsuno T, Nagata S, Sato Y, Watanabe T (1974). Ibidem. 2. Carotenoids of horse mackerel, swellfishes, porcupine fishes and striped mullet. *Bull. Jap. Soc. Sci. Fish.* 40:579-584.
- McDonald G (1995). Early mortality syndrome (EMS) in Baltic salmon backgrounds notes. *Research Workshop, Detroit, Michigan* 2-3 February 1995: 10-12.
- Miki W, Yamaguchi K, Konosu S (1982). Comparison of carotenoids in the ovaries of marine fish and shellfish. *Comp. Biochem. Physiol.* 71B:7-11.
- Palozza P, Serini S, Ameruso M, Verdecchia S (2009). Modulation of intracellular signalling pathways by carotenoids. In: Britton G et al. (Eds.), *Carotenoids*, vol. 5: Nutrition and health, Birkhäuser verlag, Basel, Boston, Berlin, pp 211-234.
- Pettersson A, Liguell A (1999). Astaxanthin deficiency in eggs and fry of Baltic salmon (*Salmo salar*) with the M74 syndrome. *Ambio* 28: 43-47.
- Pfander H, Riesen R (1995). Chromatography: Part IV. High-performance liquid chromatography. In: Britton G et al. (Eds.), *Carotenoids*, vol. 1A: Isolation and analysis, Birkhäuser verlag, Basel, Boston, Berlin, pp 145-190.
- Rock CL (2009). Carotenoids and cancer. In: Britton G et al. (Eds.), *Carotenoids*, vol. 5: Nutrition and health, Birkhäuser verlag, Basel, Boston, Berlin, pp 269-286.
- Schiedt K (1995). Chromatography. Part III. Thin- layer chromatography. In: Britton G et al. (Eds.), *Carotenoids*, vol. 1A: Isolation and analysis, Birkhäuser verlag, Basel, Boston, Berlin, pp 131-144.
- Schiedt K, Liaaen- Jensen S (1995). Isolation and analysis. In: Britton G et al. (Eds.), *Carotenoids*, vol. 1A: Isolation and analysis, Birkhäuser verlag, Basel, Boston, Berlin, pp 81-108.
- Simpson KL, Katayama T, Chichester CO (1981). Carotenoids in fish feeds. In: Baurenfeindt JC (Ed.), *Carotenoids as Colorants and Vitamin A Precursors*, vol. 4, Academic Press, New York, pp 463-538.
- Straub O (1987). *Key to Carotenoids*, Birkhäuser Verlag, Basel- Boston, 296 pp.
- Tanaka Y, Nakano F, Katayama T (1978). The carotenoids In tropical marine yellow fish. *Mem. Fac. Fish. Kagoshima Univ.* 27:29-33.
- Torrisen OJ, Hardy RW, Shearer KD (1989). Pigmentation of salmonids- carotenoid deposition and metabolism. *Crit. Rev. Aquat. Sci.* 1: 209-225.
- Tsukuda N, Amano K (1966). Studies on the discoloration of red fishes. 1. content of carotenoid pigments in eighteen species of red fishes. *Bull. Jap. Soc. Sci. Fish.* 32:334-345.
- Vetter W, Englert G, Rogassi N, Schwieter U (1971). Spectroscopic methods. In: Isler O (Ed.), *Carotenoids*, Birkhäuser Verlag, Basel- Boston, pp189-229.
- Vuorinen PJ, Paasivirta J, Keinänen M, Kostinen J, Rantio T, Hyötyläinen T, Welling L (1997). The M74 syndrome of Baltic salmon (*Salmo salar*) and organochlorine concentrations in the muscle of female salmon. *Chemosphere* 34: 1151-1166.
- Wang X-D (2009). Biological activities of carotenoid metabolites. In: Britton G et al. (Eds.), *Carotenoids*, vol. 5: Nutrition and health, Birkhäuser verlag, Basel, Boston, Berlin, pp 383-408.
- Winer BJ (1997). *Statistical principles in experimental design*. McCraw Hill, New York, 375 pp.
- Yeum K-J, Aldini G, Russell RM, Krinsky NI (2009). Antioxidant/ pro- oxidant actions of carotenoids. In: Britton G et al. (Eds.), *Carotenoids*, vol. 5: Nutrition and health, Birkhäuser verlag, Basel, Boston, Berlin, pp 235-268.