

Full Length Research Paper

Enhancing rotifer mass culture through algal diversity

Paul O. Ajah

Institute of Oceanography, University of Calabar, Calabar, Cross Rivers State, Nigeria. E-mail: ajapaulo@yahoo.com.
Tel: +2348033707901.

Accepted 4 January, 2024

Outdoor cultures of *Brachionus quadridentatus* raised in 10 m³ concrete tanks fed on co-cultured three algal species (*Chlorella vulgaris*, *Eudorina elegans* and *Scenedesmus quadricauda*) feed showed statistically significant differences in population biomass growth. Growth of *B. quadridentatus* feeding on *S. quadricauda* was significantly greater than when fed on *C. vulgaris*. No significant differences in growth were observed between *Brachionus* cultures fed on *S. quadricauda* and *E. elegans* and between those fed on *C. vulgaris* and *E. elegans*. Daily *B. quadridentatus* population biomass increase was highest with *S. quadricauda*, followed by *E. elegans* and lastly *C. vulgaris*. *B. quadridentatus* needed approximately 20, 48 and 63 h, respectively, with *C. vulgaris*, *S. quadricauda* and *E. elegans*, to double its population.

Key words: Zooplankton, algae, culture, nutrient.

INTRODUCTION

Mass production of catfish under controlled conditions depends on the provision of live plankton food for early fry and larval stages. The importance of live food in fry and larval rearing has been reported (Ovie et al., 1993; Ajah and Holzlohner, 1996; Ajah, 1997, 1998; Hagiwara et al., 1997, 2007). The successful production of finfish or shellfish such as shrimps and crabs not only requires a starter diet such as *Lepadella ovalis* (L-type (93.6 × 70.3 μm²) or S-type (70.3 × 35.1 μm²) (Ajah and Ajah, in press), but also a second diet such as *B. quadridentatus* (432.9 × 128.7 μm²) and a third diet, for example, *Asplanchna* (685 × 356 μm²) (Ajah, 2008). A number of authors have emphasized the importance of adequate supply of rotifer cultures with the appropriate nutritional quality, for the survival of larvae cultured in marine hatcheries (Hirata, 1980; Kafuku and Ikonoune, 1983; Watanabe et al., 1983; Lubzens, 1987). Previous authors like Ovie et al. (1993); Adeyemo et al. (1994) and Ajah (1997, 1998) have established the advantages of freshwater plankton over *Artemia nauplii*, a salt-water Anostraca. It is best to culture freshwater fish with freshwater live foods (Ovie et al., 1993; Ajah, 1997). *B. quadridentatus*, a rotiferan, could be a suitable alternative to *A. nauplii*, being of equivalent size to imported *A. nauplii*. Another reason for preferring *B. quadridentatus* is that, it occurred quite frequently and in large numbers in ponds at the University of Calabar fish farm. Earlier, Lubzens et

al. (1990) stated that rotifers (*Brachionus plicatilis*) constitute a major, and in most cases the only food source for larval stages of several organisms raised in marine aquaculture including fish and various invertebrates.

The culture of at least 10 species of marine rotifers including *B. plicatilis*, *Brachionus rotundiformis*, *Brachionus pterodinooides*, *Brachionus satanicus* and *Hexarthra jenkiniae* is well documented (Gatesoupe and Luquet 1981; Lubzens, 1987; Lubzens et al., 1989, 1990; Hampton and Starkweather, 1998). Gatesoupe and Luquet (1981) noted that there could be nutritional differences in the value of rotifers as food for larvae based on the feeding conditions of the rotifers themselves. Watanabe et al. (1979), found that *Chlorella*-fed and yeast-fed rotifers differed in their proximate compositions. Ajah (1998) found increased growth and survivorship of *Heterobranchus longifilis* larvae when fed on enriched zooplankton.

Earlier, Fujita (1979) indicated the importance of long chain ω-3 polyunsaturated fatty acids in rotifers as food for Red Sea bream larvae, and found a further improvement in the dietary value of yeast cultured rotifers by secondary culture with marine *Chlorella* for 6 h. *Senecio quadridentatus* has been found to thrive better in freshwater cultures than *B. plicatilis* (Ajah, 1995).

One advantage of *B. quadridentatus* over *B. plicatilis* is its low failure rate during culture. *B. quadridentatus*

culture was maintained continuously for two years irrespective of seasons with little fluctuation in population size (Ajah, 1995). The main difference between the dry and wet season blooms is in the initiation of blooms which brought about a doubling of the initial nutrients for the first two days during the wet season (Ajah, 1995).

Brachionus calyciflorus, a freshwater rotifer was successfully fed on two types of algae, *Scenedesmus obliquus* and *Chlorella* sp. (Kennari et al., 2008).

In this study, freshwater food items (*Chlorella vulgaris*, *Eudorina elegans* and *Scenedesmus quadricauda*) that might be more suitable, acceptable, cheap, readily available and capable of yielding higher population densities of *B. quadridentatus* a freshwater rotifera were investigated.

MATERIALS AND METHODS

Three hard bottomed 10 m³ circular concrete tanks with central drainage pipes and five airlift pipes, arranged in an anticlockwise pattern to keep the plankton culture afloat were used for the outdoor culture of the rotifer. Axenic monocultures of micro algae (*C. vulgaris* Beij. var. *vulgaris* Fott, (2 - 12 µm), *S. quadricauda* (Turp.) Bréb (12 - 15 µm) and *E. elegans* Ehr. (95 - 105 µm) as well as pure monocultures of *B. quadridentatus* (Hermann, 1783), were cultured in the laboratory of the Hatchery Complex of the Institute of Oceanography, University of Calabar, Calabar, Nigeria, using both the "Gelose" and "Dilution" isolation methods (Harder, 1917; McVey and Moore, 1983). The algae samples were collected from the fish ponds of the Institute using a plastic container and inoculated into the algal medium using Pasteur pipette. Axenic algal cultures were produced under highly hygienic laboratory conditions using autoclaving and inhibitory bacterial growth precursors such as 0.005 - 0.01% hypochlorite solution, UV radiation and antibiotics (Ajah, 1995). Species were identified using Ward and Whipple (1959), Bold and Wynne (1978) and Jeje and Fernando (1986).

Zooplankton culture

Two 36-watt daylight fluorescent tubes were suspended 60 cm above a series of 5 - 60 L aquaria to illuminate a wooden cupboard of 120 × 60 × 80 cm³ (0.6 m³). Room temperature was kept at 28 ± 1°C and aeration using two 1.5 hp pumps was continuous. The pure zooplankton cultures received 50 mg l⁻¹ of baker's yeast (*Saccharomyces cerevisiae*) or 1.75 mg l⁻¹ of inorganic fertilizer (N: P: K) (20:10:10) every other day to release the macronutrients needed for zooplankton growth, whereas 1.0 ml of nutrient salt solution A and 0.1 ml of solution C (Laing and Ayala, 1990) were administered every other day to the pure algal cultures (Ajah, 1995).

Outdoor cultures

Fertilization of outdoor cultures was made possible using pig manure supplied regularly by 20 weaned pigs that were fed daily on 10% body weight with pig mash (Akpan and Okafor, 1997). The chemical composition of samples of pig manure was determined and found to contain the essential nutrient elements for algal growth. One (1.0) kg of pig manure was introduced into each outdoor tank during the dry season and 2 kg during the wet season for the first two consecutive days. Thereafter, 1 kg per tank was administered every three days and every other day, for the dry and

the wet seasons, respectively. 50 ml of each axenic algal monoculture, namely, *C. vulgaris*, *S. quadricauda* and *E. elegans*, from the laboratory, were inoculated into the respective tanks at cell densities of 2.6 × 10⁶, 1.68 × 10⁶ and 0.12 × 10⁶ cells/ml. Five litres each of 4 ind ml⁻¹ from pure *B. quadridentatus* cultures (Ajah, 1995) were introduced into each 10 m³ tank. The feeding duration of *B. quadridentatus* with each of the three algae lasted for an average of 18 days (range 15 - 21 days).

Each culture was repeated three times during both the dry season (November to March/ April) and wet months (May/ June to October). The batch culture trials continued for a period of three years amounting to a total of eighteen trials, nine per season (Ajah, 1995, 1997, 1998).

Cell counts

A 1 l plastic funnel was used to scoop samples from the thoroughly mixed plankton cultures. The samples filtered through a 56 µm-plankton sieve to collect and retain both phyto- and zooplankton. Four cell counts of 1 ml each using a haemocytometer and four zooplankton counts using a one ml counting chamber (model: AJAH001) (Ajah, 1995) were carried out every day or every two days, and average values were recorded.

Physicochemical parameters

The physical and chemical factors were monitored throughout the experimental period. Dissolved oxygen in the pond was determined using Lectron 5509 DO meter and temperature was read using standard thermometer. Chlorine was assessed by the chlorosity method of Rump and Krist (1988). Nitrite (NO₂⁻-N) by the diazotization (spectrophotometric) method; nitrate (NO₃⁻-N) by the cadmium reduction/ diazotization method, ammonium (NH₄⁺-N) level by the Nesslerization (spectrophotometric) method, phosphate by the molybdenum blue method (spectrophotometric) (Parsons et al., 1984), conductivity was read using a HACH 3000 spectrophotometer and turbidity by secchi disc.

Single classification analysis of variance (ANOVA) was used following Sokal and Rohlf (1981) to compare means of population density of zooplankton counts from the replicates. Correlation coefficients between *B. quadridentatus* and each primary producer as well as the coefficients of determination were also calculated using SPSS. Descriptive statistics were used to compare the means. The intrinsic rate of natural increase (*r*) and the population doubling time in days (*t_D*) were calculated as follows:

$$r = (\ln N_t - \ln N_0) / t = D'/t \text{ or } 2.3026(\log N_t - N_0)/t \text{ (James and Dias, 1984).}$$

Where:

D' = The dilution time in days.

$N_t = N_0 \cdot e^{rt}$ (James and Dias, 1984).

N_t = Final number of individuals.

N₀ = Initial number of individuals, and *t* = time in days

e (exponent) = 2.7183.

r = The intrinsic rate of natural increase.

t_D = 0.6931/*r* (James and Dias, 1984).

t_D = Doubling time of the population in days.

RESULTS

C. vulgaris, *E. elegans* and *S. quadricauda* showed statistically significant ($F_{2,115} = 3.27$, $p < 0.05$) differences in

Table 1. Daily mean values and summary of the three replicates showing some growth characteristics of *Brachionus quadridentatus* while feeding on the three algae.

<i>Chlorella vulgaris</i>			<i>Scenedesmus quadricauda</i>			<i>Eudorina elegans</i>			
Date	r	t _D (time in days)	N _t (ind/L)	r	t _D (time in days)	N _t (ind/L)	r	t _D (time in days)	N _t (ind/L)
1									
4	0.207	3.508	3,000	0.414	1.801	4,000	0.496	1.435	3,000
5	0.433	1.611	15,000	0.504	1.423	18,000	0.638	1.094	8,000
8	0.326	2.129	31,000	0.381	1.835	41,000	0.371	1.369	11,000
9	0.312	2.368	50,000	0.424	1.637	65,000	0.433	1.071	32,000
10	0.284	3.484	62,000	0.291	2.414	70,000	0.286	1.615	46,000
11	0.375	1.848	123,000	0.360	1.936	153,000	0.272	2.422	93,000
12	0.373	1.860	143,000	0.354	1.948	160,000	0.096	5.722	104,000
15	0.310	2.234	124,000	0.158	4.888	155,000	0.119	5.802	118,000
16	0.279	2.488	88,000	0.207	3.366	125,000	0.319	2.169	127,000
17	0.257	2.707	63,000	0.140	4.978	94,000	0.189	3.660	98,000
19	0.198	3.500	55,000	0.155	4.472	77,000	0.170	4.111	55,000
22	0.216	0.482	25,000	0.108	9.02	46,000	0.123	5.724	36,000
Biomass indiv. × 10 ⁶ /10 m ³ /day		366.92		489.52		289.05			
Peak densities(ind./L)		102,000		104,000		176,000			
Av. pop. (N _t) (ind/day)		65,167 ± 45622		72,000 ± 51464		60,916 ± 44971			
Av. rate of natural increase (r) l/day		0.149 ± 0.01		0.276 ± 0.01		0.286 ± 0.01			
Av. Doubling time of pop. (t _D) in days.		0.84 ± 0.89		2.04 ± 0.77		2.61 ± 0.38			
R ² b/w <i>Brachionus</i> and alga density		0.166, p < 0.05		0.044, p > 0.05		0.162, p > 0.05			

population biomass growth and development. *B. quadridentatus*' growth using *S. quadricauda* was significantly (¹,₇₆F6.804, P 0.01) greater than when fed on *C. vulgaris*. There were no significant (¹,₇₇F0.330, P 0.05; ¹,₇₆F3.499, P 0.05, respectively) growth differences between *B. quadridentatus*-fed, *S. quadricauda* and *E. elegans*, and, between *B. quadridentatus* fed on *C. vulgaris* and *E. elegans*. Correlation coefficients between *B. quadridentatus* densities fed on *C. vulgaris* and *E. elegans*, *E. elegans* and *S. quadricauda* and *S. quadricauda* and *E. elegans* were 0.454, 0.413 and 0.618, respectively.

The population biomass of *B. quadridentatus* increased quite rapidly with *S. quadricauda*, followed by *E. elegans* and lastly *C. vulgaris* (Table 1). High average population densities of *B. quadridentatus* achieved with *C. vulgaris*, *E. elegans* and *S. quadricauda* are shown in Table 1. *B. quadridentatus* showed extremely high rates of natural increase, with correspondingly very short population doubling times (t_D) with *S. quadricauda* (Table 1). The intrinsic rate of natural increase (r) and doubling time in days (t_D) for *B. quadridentatus* using *E. elegans*, and *C. vulgaris* cultures are equally represented in Table 1.

The overall daily mean population growth (N_t)

was 65,167 ± 45622, 60,916 ± 44971 and 72,000 ± 51464 for *C. vulgaris*, *E. elegans* and *S. quadricauda*, respectively. However, based on the three replicates, *B. quadridentatus* needed less time, on the average 0.84 days, to double its population when fed on *C. vulgaris*, followed by 2.04 and 2.61 days, respectively, was required by *S. quadricauda* and *E. elegans* to double their populations (Table 1). More stable and higher growth rates were obtained using *S. quadricauda* and *E. elegans* compared to *C. vulgaris*. The chemical composition of the swine manure was: NH₄⁺-N = 178.3 mg l⁻¹, SO₄ = 85.44 mg l⁻¹, SiO₂ = 85.2 mg l⁻¹, NO₂⁻-N (below detectable level), NO₃⁻.




Figure 1. Showing seasonal mean physicochemical factors and nutrient contents obtained during the culture of *Brachionus quadridentatus* in earth pond.

N = 25.3 mg l^{-1} , PO $_4^{3--P}$ = 0.64 mg l^{-1} , N = 28.7 ppm/22.7%, P = 0.20 ppm/0.66%, K = 12.5 ppm/31.96% and moisture content 62.6%. The percentages represent total available nitrogen, phosphorus, potassium and moisture content in the sample. The mean values and ranges of the physicochemical parameters and nutrient loads of the culture environments are shown in Figure 1. Temperature ranged from 26.5°C to 29.4°C with average of $27.74 \pm 0.108^\circ\text{C}$. Dissolved oxygen levels ranged from 2.7 - 5.4 mg l^{-1} with a mean of $4.238 \pm 0.126 \text{ mg}l^{-1}$. Mean conductivity values in the culture tank were $642.567 \pm 142 \text{ s/cm}$ ranging from 153.7 - 921 s/cm. The turbidity of the culture system ranged from 109 - 279 FTU with a mean of $218.607 \pm 32 \text{ FTU}$. The NO $_3$ -N level range from 1.081 to 3.050 mg l^{-1} with a mean of $2.048 \pm 0.549 \text{ mg}l^{-1}$ while the phosphate levels in the tanks were 0.150 to 1.016 mg l^{-1} with a mean of $0.647 \pm 0.149 \text{ mg}l^{-1}$.

DISCUSSION

The physicochemical factors and nutrient load were within the acceptable limits required for the proper management of earth ponds (Wedemeyer et al., 1976; Post, 1987; Swift, 1988; Agarwal, 1994; Biswas, 1996; Ajah, 2008). Physicochemical values in this report that fell within the optima required for plankton growth included amongst others temperature (IFAS Circular, 1951), DO (Banerjee, 1967), the electrolyte conductivity, pH and the DO (Ajah, 1995, 2008), turbidity (Ajah, 1995, 2008), nitrate (Sachidanandamurthy and Yajurvedi, 2004) and phosphate (Sawyer, 1947; Screenivasan, 1965).

B. quadridentatus due to its extremely high fecundity doubled its population in only 0.149 days and had a t_D of 0.84, giving it a comparative advantage over *B. plicatilis* that had $r = 0.84$, $t_D = 1.212$ obtained with

Nannochloropsis sp. (Ahmad et al., 1991). This puts *B. quadridentatus* as a more promising alternative food source to *Artemia*.

The greater preference for *S. quadricauda* and *E. elegans* to *C. vulgaris* by *B. quadridentatus* could be explained in terms of the voracious nature and size of *Brachionus* as well as its choice of food item. Kennari et al. (2008) rather observed that *B. calyciflorus* had better growth and fatty acid content when fed on *Chlorella* sp. compared with *S. obliquus*. Investigations revealed that phytoplankton abundance was controlled by zooplankton abundance and not fertilization as in a hard water pond (Sierp, 2001), whereas in soft bottom ponds such as ours both zooplankton abundance and fertilization controlled phytoplankton abundance. Ajah (2008) earlier reported a greater preference of *Asplanchna priodonta* for *E. elegans* which confirms the need for larger prey by larger predators.

The initial population growth of *Keratella cochlearis* was significantly affected by the feeding schedule as well as the presence of competitors, while that of *Daphnia* was affected by neither factor. Population densities of both species tended to increase as the frequency of food addition increased (Maclsaac and Gilbert, 1991). Xi et al. (2002) found out variations in population growth, body size and egg size of *B. calyciflorus* while feeding on different algae. Fabiola et al. (2005) equally observed differences in population growth of rotifers and cladocerans when raised on algal diets supplemented with yeast. Mean population growth rate (r) of 0.61 and 0.44 were obtained for rotifer fed with *Chlorella* sp. and *S. obliquus*, respectively (kennari et al., 2008) which are lower than values obtained in this report using three algae, *C. vulgaris* (0.149), *S. quadricauda* (0.276) and *E. elegans* (0.286).

Conover and Mayzaud (1984) made the following

generalizations regarding selective grazing: (1) At low concentrations of food, feeding tends to be spread over a wide range of particle sizes, (2) Although “tracking” or selection of biomass peaks as a strategy to facilitate capture of rapidly reproducing, and presumably more nutritious phytoplankton (Poulet and Chanut, 1975), and (3) If the particle has a “suitable” shape and is not chemically distasteful, only then do zooplankton tend to select bigger objects. Points 2 and 3 above agree with the findings made here. My investigation showed that there was an almost equal attraction for the various food items administered, proving that none of the food items had unpleasant odour or taste to *B. quadridentatus*, nor did any possess an awkward shape.

In conclusion, prey size, shape, and motion are implicated as probable causes for the slight differences in feeding behaviour. *B. quadridentatus* generally preferred the bigger prey like *E. elegans*, or medium size prey, but with some flagella-like structure as in *S. quadricauda* as well as prey in constant motion such as *E. elegans*.

The cilia of *B. quadridentatus* were capable of generating waves of current for trapping its prey. Distance was not a barrier for capturing prey, due to its fast motion. *B. quadridentatus* could easily be grown on three algal live diets such as *C. vulgaris*, *E. elegans* and *S. quadricauda*.

ACKNOWLEDGEMENT

This study was funded through a fellowship provided by the European Economic Community.

REFERENCES

- Adeyemo AA, Oladosu GA, Ayinla AO (1994). Growth and survival of African catfish species, *Claria gariepinus* Burchell, *Heterobranchus bidorsalis* Geoffrey and *Heteroclinas* reared on *Moina dubia* in comparison with other first feed sources. *Aqua*. 119: 41-45
- Agarwal SC (1994). A Handbook of Fish Farming. Narendra Publishing House, India p. 117
- Ahmad TA, Yamasaki S, Hirata H (1991). Optimum feeding rate of the rotifer *Brachionus plicatilis* on the marine alga *Nannochloropsis* sp. *J. World Aquacult. Soc.* 22: 230-234.
- Ajah PO (1995). Identification and mass production of zooplankton species suitable for early feeding of African clariid catfish. PhD thesis, University of Calabar, Nigeria p.164.
- Ajah PO (1997). Effects of live foods, artificial feed and their combination on growth and survival of African clariid catfish (*Heterobranchus longifilis* Valenciennes, 1840) larvae. *The Israeli J. Aquacult.* 49: 205-213.
- Ajah PO (1998). A comparison of growth and survival of *Heterobranchus longifilis* fed on *Artemia nauplii* and nine non-*Artemia* live diets. *Trop. Freshwater Biol.* 7: 1-15.
- Ajah PO, Ajah MO (in press). Comparative Growth Performance of Freshwater Zooplankton (*Lepadella ovalis* Borys de St. Vincent, a Rotifer) on three algal species. *J. Plankton Res.*
- Ajah PO (2008). Growth Characteristics of the Monogonont Rotifer *Asplanchna priodonta* Gosse 1850 on Three Algae Species. *Turk. J. Fish. Aqua. Sci.* 8: 275-282.
- Ajah PO, Holzlohner S (1996). The use of low cost organic manure in mass-producing zooplankton. In: Future trend in aquaculture development in Eastern Europe. Handbook of short communications and National Reports, Budapest, Hungary pp. 3-4.
- Akpan ER, Okafor N (1997). On organic fertilization and plankton development in two experimental freshwater ponds of Nigeria. *J. Aquacult. Trop.* 12: 147-154.
- Banerjea SM (1967). Water quality and soil condition of fish ponds in some states of India in relation to fish production. *Indian J. Fish.* 14: 115-144.
- Biswas KP (1996). A Text Book of Fish, Fisheries and Technology. Narendra Publishing House. India p. 578.
- Bold HC, Wynne MJ (1978). Introduction to the algae: Structure and Reproduction. Bounty...Prentice-Hall Inc., Englewood Cliffs, New Jersey p. 707.
- Conover RJ, Mayzaud P (1984). Utilization of Phytoplankton by Zooplankton during the Spring Bloom in a Nova Scotia Inlet. *Canadian J. Fish. Aqua. Sc.* 41: 232-244.
- Fabiola PA, Nandini S, Sarma SSS (2005). Differences in population growth of rotifers and cladocerans raised on algal diets supplemented with yeast. *Limnologia-Ecology and Management of Inland Waters* 35(4): 298-303.
- Fujita S (1979). Culture of Red Sea Bream, *Pagrus major*, and its food. In: European Mariculture Society Special Publication 4: 183-197.
- Gatesoupe FJ, Luquet P (1981). Practical diet for mass culture of the rotifer *Brachionus plicatilis*: application on larval rearing of sea bass, *Dicentrarchus labrax*. *Aquacult.* 22: 149-163.
- Hagiwara A, Snell TW, Lubzens E, Tamaru CS (1997). 'Live food in aquaculture' *Developments in Hydrobiology* 124, Kluwer: p. 328.
- Hagiwara A, Suga K, Akazawa A, Kotani T, Sakakura Y (2007). Development of rotifer strains with useful traits for rearing fish larvae. *Aquacult.* ISSN 0044-8486 CODEN AQCAL.
- Hampton SE, Starkweather PL (1998). Differences in predation among morphotypes of the rotifer *Asplanchna silvestrii*. *Freshwater Biology*, 40(4): 595 -doi:10.1046/j.1365-2427.1998.00359.x.
- Harder R (1917). Emhrungsphysiologische Untersuchungen an Cyanophyceen, hauptsächlich dem endophytischen *Nostoc punctiforme*. *Z. Botanical* 9: 145-242.
- Hirata H (1980). Culture methods of marine rotifer *Brachionus plicatilis*. *Min. Rev. Data File Fish. Res.* 1: 27-46.
- IFAS, Circular 1051. Institute of food and agriculture sciences, University of Florida.
- James CM, Dias P (1984). Mass culture and production of the rotifer *Brachionus plicatilis* using baker's yeast and Marine yeast. Annual Research Report Kuwait Institute for Scientific Research pp. 49-51.
- Jeje CY, Fernando CF (1986). A practical guide to the identification of Nigerian zooplankton (Cladocera, Copepoda and Rotifera). Published by Kainji Lake Research Institute, New Bussa p. 742.
- Kafuku T, Ikonoune H (1983). Modern methods of aquaculture in Japan. *Developments in Aquaculture and Fisheries Science*, 11. Kodansha Ltd., Tokyo, and Elsevier, Amsterdam p. 216.
- Kennari AA, Ahmadifard N, Sevfabadi J, Kapourchali MF (2008). Comparison of growth and fatty acids composition of freshwater Rotifer, *Brachionus calyciflorus* Pallas, fed with two types of microalgae at different concentrations. *J. World Aquacult. Soc.* 39(2): 235-242.
- Laing I, Ayala F (1990). Commercial mass culture techniques for producing microalgae. *Introduction to Applied Phycology. Archive Hydrobiologia/Supplement* 5: 64-67.
- Lubzens E (1987). Raising rotifers in aquaculture (a review). 1V. Rotifer Symposium, Edinburgh, Scotland. *Hydrobiologia* 147: 245-255.
- Lubzens E, Tandler A, Minkoff G (1989). Rotifers as food in aquaculture. *Hydrobiologia* 186/187: 387-400.
- Lubzens E, Kolodny G, Perry B, Galai N, Sheshinski R, Wax Y (1990). Factors affecting survival of rotifers (*Brachionus plicatilis* O.F. Müller) at 4°C. *Aquacult.* 91: 23-47.
- MacIsaac HJ, Gilbert JJ (1991). Competition between *Keratella cochlearis* and *Daphnia ambigua*: effects of temporal patterns of food supply. *Freshwater Biol.* 25: 189-198.
- Mayeli SM, Nandini S, Sarma SSS (2004). The efficacy of *Scenedesmus* morphology as a defense mechanism against grazing by selected species of rotifers and cladocerans. *Aqua. Ecol.* 38(4): 515-524.
- McVey JP, Moore JR (1983). Hatchery techniques for Penaeid species fed six food combinations. *Aquacult.* 47: 151-162.

- Mohr S, Adrian R (2002). Reproductive success of the rotifer *Brachionus calyciflorus* feeding on ciliates and flagellates of different trophic modes. *Freshwater Biol.* 47(10): 1832 - doi:10.1046/j.1365-2427.2002.00929.x
- Morales-Ventura J, Nandini S, Sarma SSS (2004). Functional responses during the early larval stages of the charal fish *Chirostoma riojai* (Pisces: Atherinidae) fed rotifers and cladocerans. *Appl. Ichthyol.* 20(5): 417 -, doi:10.1111/j.1439-0426.2004.00565.x.
- Nandini S, Pérez-Chávez R, Sarma SSS (2003). The effect of prey morphology on the feeding behaviour and population growth of the predatory rotifer *Asplanchna sieboldi*: a case study using five species of *Brachionus* (Rotifera). *Freshwater Biol.* 48 (12): 2131 - doi:10.1046/j.1365-2427.2003.01149.x.
- Nandini S, Sarma SSS (1999). Effect of starvation time on the prey capture behaviour, functional response and population growth of *Asplanchna sieboldi* (Rotifera). *Freshwater Biol.* 42 (1): 121 -doi: 1046/j.1365-2427.1999.00467.x.
- Ovie SI, Adeniji HA, Olowe DI (1993). Isolation and growth characteristics of a freshwater zooplankton for feeding early larval and fry stages of fish. *Aquacult. Trop.* 8: 187-196 .
- Post G (1987). *Text Book of Fish Health*. T.F.H Publications, Inc. USA.
- Poulet SA, Chanut JP (1975). Non-selective feeding of *Pseudocalanus minutus*. *J. Fish. Res. Board Canada* 32: 706-713.
- Sachidanandamurthy KL, Yajurvedi HN (2004). Monthly variations of water quality parameters (Physico-chemical) of a perennial Lake in Mysore City. *Indian Hydrobiol.* 7: 217-228
- Sawyer CN (1947). Fertilization of lake by agriculture and urban drainage. *J. New Water Lakes Ass.* 51: 109-127.
- Screenivasan A (1965). Limnology and productivity of tropical upland impoundment s in Nilgiris, Madras state. *India Phytos.* 7: 146-160.
- Sierp MT (2001). Effects of fertiliser and crayfish on plankton and nutrient dynamics in hard water ponds. *Hydrobiologia* 462: 1-7.
- Sokal RR, Rohlf FJ (1981). *Biometry. The principles and practices of statistics in biological research.* 3rd Ed New York. Freeman and Company.
- Swift DR (1988). *A manual of aquaculture training.* Fishing News Books Ltd. England p 135.
- Ward BH, Whipple CG (1959). *Freshwater Biology.* 2nd ed. W. T. Edmondson ed. John Wiley and Sons, Inc. New York p.1248.
- Watanabe T, Oowa F, Kitajima C, Fujita S, Yone Y (1979). Relationship between the dietary values of *Brachionus plicatilis* and their content of ω -3 highly unsaturated fatty acids. *Bull. Jpn Soc. Sci. Fish* 45: 883-889.
- Watanabe T, Kitajima C, Fugita S (1983). Nutritional values of live organisms used in Japan for mass propagation of fish. *Rev. Aquacult.* 34: 115-143.
- Wedemeyer GA, Meyer FP, Smith L (1976). *Diseases of Fishes: Environmental stress and fish diseases* pp. 89-106.
- Xi YLI, Liu GYI, Jin HJ (2002). Population growth, body size, and egg size of two different strains of *Brachionus calyciflorus* Pallas (Rotifera) fed algae. *J. Freshwater Ecol.* 17(2): 185-190.

