

Full Length Research Paper

ST Segment Elevation in Leads V5 and V6 for Predicting Culprit Artery in Acute Inferior Wall Myocardial Infarction

Madhu Gupta¹, Maheshwar Prasad², Sun Jian¹, Rajesh Panjiyar³, Shankar Laudari^{3*} and G Subramanyam³

¹Cardiovascular Disease Center, The first hospital of Jilin University, Norman Bethune College of Medicine, Changchun, PR China, 130021

²Department of Medicine, National Medical College, Nepal

³Department of Cardiology, College of Medical Science, Nepal

Accepted 15 September, 2022

ST elevation (ST \uparrow) in the leads V5 and V6 to predict the culprit and furthermore, comparing the degree of ST \uparrow in lead V6 with that in lead III. Patients were first divided according to the presence (n-62) and absence (n-98) with ST \uparrow of ≥ 2 mm in leads V5 and V6, the patients were then further subdivided into the 2 groups according to the degree of ST \uparrow in lead III and in lead V6 and were compared with coronary angiography. In patients with ST \uparrow in leads V5 and V6, the infarct related artery was right coronary artery (RCA) in 66% and left circumflex artery (LCx) in 31%. The RCA was found significantly higher proportion in ST \uparrow in leads V5 and V6 with ST \uparrow in lead III $>$ V6 (n-37; 82%) ($p=0.0001$), whereas the LCx disease in ST \uparrow in leads V5 and V6 with ST \uparrow in III \leq V6 (n-12; 71%) ($p=0.0001$). The sensitivity, specificity, positive and negative predictive value in ST \uparrow in leads V5 and V6 with ST \uparrow in lead III $>$ V6 for RCA, and ST \uparrow in lead III \leq V6 for LCx to predict culprit artery were 90%, 63%, 84%, 75% and 63%, 90%, 75%, 84% respectively. ST elevation in leads V5 and V6 suggests a greater risk area in patient with acute inferior wall myocardial infarction. It also signifies a larger perfusion territory and demands for more aggressive reperfusion therapy.

Keywords: Infarct related artery (IRA); ST elevation myocardial infarction (STEMI); Right coronary artery (RCA); Left circumflex artery (LCX); Electrocardiogram (ECG).

INTRODUCTION

The inferior wall and inferolateral wall of the left ventricle are supplied by the right coronary artery (RCA) and left circumflex (LCx) artery. Inferior wall acute myocardial infarction (I-AMI) is caused by an occlusion of either the

RCA or LCx. Identification of RCA occlusion is very important because it is associated with the right ventricle extension and the clinical syndrome of hypotension and shock (Braat Set al., 1984). Multiple ECG criteria have been studied to predict the culprit artery based on ST segment elevation and ST segment depression in different leads (Kosuge et al., 1998; Chia et al., 2000; Herz et al., 1997; Zimetbaum et al., 1998; Bairey et al., 1987; Hasdai et al., 1995; Assali et al., 1999; Fiol et al.,

*Corresponding Author E-mail: Ishankar2@hotmail.com;
Phone: 9845112909

2004; Fiol et al., 2004). The present study examines an electrocardiographic algorithm based on ST segment changes in leads V5 and V6 that has a very high sensitivity, specificity and predictive value to predict the culprit artery in an inferolateral acute myocardial infarction.

METHODS

The present study was approved by our local hospital Institutional Review Board (IRB). We retrospectively analyzed 390 patients having an acute inferior wall MI, admitted and underwent coronary angiography in the cardiovascular department of our hospital within a year-period. Only 160 patients (male 132 and female 28) met our inclusion criteria of chest pain for ≥ 30 minutes before hospital admission. If the elevation of creatinine kinase (CK-MB) is greater than twice the upper limit (Normal: 0-3.5 ng/ml), the ECG shows ST segment ≥ 0.1 mV (1mm) in at least 2 of 3 inferior leads (II, III, aVF). Coronary angiography showing total occlusion or critical stenosis $>70\%$ in single vessel either RCA or LCX. Whereas, exclusion criteria include lack of ST elevation ≥ 0.1 mV (1mm) in the inferior leads (II, III, aVF), previous history of acute myocardial infarction, coronary artery bypass surgery or percutaneous coronary intervention prior to current hospitalization, evidence of recent left bundle branch block or left ventricular hypertrophy on ECG, and significant stenosis in both LCX and RCA or triple vessel disease so that a single infarct related artery could not be defined. A 12 leads ECG was recorded in all patients at a paper speed of 25mm/s and voltage 10mm/mV. The ST-elevation deviation was measured manually to the nearest 0.5mm, 80ms after the J point by a single cardiologist who was unaware of all clinical and angiographic findings. ST segment depression in aVR of at least 0.1mV (1mm) was the most important criterion to distinguish between LCx and RCA as presumptive predictor of the culprit artery. Coronary angiography was performed during hospitalization in all patients. The results of this test were compared with those obtained from ECG to distinguish the real culprit artery. Data are expressed as the mean \pm SD for continuous variable and percentages for categorical variables. Statically significance ($p < 0.05$) was assessed by chi-square test or fisher exact test for categorical variables. The sensitivity, specificity, positive predictive value and negative predictive value were evaluated in the 2 coronary arteries.

RESULTS

A total of 160 patients (male 132 and female 28) was included in the study. Patients were divided according to the presence (n -62) and absence (n -98) of ST elevation of ≥ 2 mm in leads V5 and V6 on admission. Patients with ST elevation in leads V5 and V6 were further subdivided into the 2 groups according to the degree of ST elevation in lead III and in lead V6. Patients with ST elevation in lead III greater than in V6 were 45 (73%), and ST elevation in lead III equal to or less than in V6 were 17 (27%) as shown in the Figure 1.2. The baseline characteristics of groups are summarized in the Table 1.1. There were no significant differences between 3 groups in baseline clinical characteristics and prevalence of risk factors. ST elevation in leads II, III and aVF with ST elevation in leads V5 and V6 was more common in male than in female (81% vs 19%).

The clinical, electrography and echocardiography findings on admission are listed in the Table 1.2. ST elevation in leads V5 and V6 with ST elevation in lead III $> V6$ was associated with greater ST elevation in leads II, III and aVF, in particular lead III. In the ECG of ST elevation in inferior leads (II, III and aVF), ST elevation in lead III $> II$ was frequently associated with ST elevation in leads V5 and V6 and ST elevation in lead III $> V6$ than in the ST elevation in leads V5 and V6 with ST elevation in lead III $\leq V6$ (84% vs 53%) ($P=0.027$), whereas ST elevation in lead II $> III$ was more associated with ST elevation in leads V5 and V6 with ST elevation in lead III $\leq V6$ than in the ST elevation in V5 and V6 with ST elevation in lead III $> V6$ (46% vs 15%) ($P=0.027$) as shown in the Table 1.2. However, ST segment depression in V1-V3 was greater in both patients with ST elevation in leads V5 and V6 with ST elevation in lead III $> V6$ and ST elevation in leads V5 and V6 with ST elevation in lead III $\leq V6$.

There was no much difference in creatinine kinase (CK-MB) and ejection fraction between three groups. Coronary angiography findings are listed in the Table 1.3. Among 62 patients with ST elevation in leads V5 and V6, the infarct related artery was right coronary artery in 41 patients (66%) and left circumflex artery in 19 (31%). No ST elevation in leads V5 and V6 and ST elevation in leads V5 and V6 with ST elevation in lead III $> V6$ groups were strongly associated with RCA occlusion; 73% and 82% respectively. However, ST elevation in leads V5 and V6 with ST elevation in lead III $\leq V6$ was associated with LCX; 71% occlusion. TIMI 0 coronary flow was found in 145 patients (91%). No differences were found between




Figure 1.1 (A). Electrocardiogram of a patient with Inferior wall acute myocardial infarction with ST-segment elevation in leads II, III and aVF and in precordial leads V5 and V6 (≥ 2 mm). ST segment elevation in leads V5 and V6 with ST elevation in lead III greater than V6. The coronary angiography later confirmed right coronary artery. (B) Electrocardiogram of a patient with Inferior wall acute myocardial infarction with no ST-segment elevation in precordial leads V5 and V6.




Figure 1.2. Shows no. of patients in inferior wall myocardial infarction with and without ST segment elevation in leads V5-V6.

Table 1.2. Clinical, Biochemical, ECG and ECHO

Variable	No ST ↑ in Lead V5-V6 (n-98)	ST ↑ in lead V5-V6		p value
		ST ↑ in lead III>V6 (n-45)	ST ↑ in lead III ≤ V6 (n-17)	
KILIP 1	93 (94%)	41 (91%)	17 (100%)	0.37
ST ↑ in				
II>III	21 (21%)	7 (15%)	8 (47%)	0.027*
III>II	77 (79%)	38 (84%)	9 (53%)	0.027*
ST in lead V1-V3				
ST depression	36(37%)	32 (71%)	12(70%)	0.0001***
Ejection Fraction	55.78±7.33	54.15±6.41	54.35±6.24	0.091
CKMB	42.26±40.69	52.43±33.05	59.35±27.61	0.178

Table 1.3. Coronary angiography (CAG)

Variable	No ST ↑ in Lead V5-V6 (n-98)	ST ↑ in lead V5-V6		P value
		ST ↑ in lead III>V6 (n-45)	ST ↑ in lead III ≤ V6 (n-17)	
RCA	72 (73%)	37 (82%)	4 (23%)	<0.0001***
LCX	22 (22%)	7 (16%)	12 (71%)	< 0.0001***
MULTIVESSELS	4 (4%)	1 (2%)	1 (5%)	0.76
TIMI 0	87 (89%)	41 (91%)	17 (100%)	0.33

Table 1.4. According to Coronary angiography; to predict the culprit artery in ST ↑ in lead

V5-V6 in related to ST ↑ in lead III>V6 and ST ↑ in lead III ≤ V6 (RCA or LCX):

Sensitivity, Specificity, PPV and NPV

ECG	CAG	Sensitivity	Specificity	PPV	NPV
ST ↑ in lead III>V6	RCA	90%	63%	84%	75%
ST ↑ in lead III ≤ V6	LCX	63%	90%	75%	84%

PPV-Positive Predictive Value, NPV-Negative Predictive Value, CAG-Coronary angiography.

three groups regarding TIMI 0 flow score. Multivessels disease were not significant.

The right coronary artery (RCA) disease was found significantly higher proportion in ST elevation in leads V5 and V6 with ST elevation in lead III > V6 (n-37; 82%) than in ST elevation in leads V5 and V6 with ST elevation in lead III ≤ V6 (n-4; 23%) ($p=0.0001$), whereas the left circumflex artery (LCX) disease was found very frequently in ST elevation in leads V5 and V6 with ST elevation in III ≤ V6 (n-12; 71%) than in ST elevation in leads V5 and V6 with ST elevation in lead III > V6 (n-7; 16%) ($p=0.0001$).

According to coronary angiography findings the sensitivity, specificity and positive predictive value and negative predictive value in ST elevation in leads V5 and V6 with ST elevation in lead III>V6 to predict right coronary artery (RCA) as a culprit artery were 90%, 63%, 84% and 75% respectively. And, the sensitivity, specificity, positive predictive value and negative predictive value in ST elevation in leads V5 and V6 with ST elevation in lead III≤V6 to predict left circumflex artery (LCX) as a culprit artery were 63%, 90%, 75% and 84% respectively as shown in the Table 1.4.

DISCUSSION

Abnormalities on the electrocardiogram (ECG) are frequently used to predict the infarct related artery (IRA) in patients with acute myocardial infarction (AMI). Acute myocardial infarction of the inferolateral wall is due to occlusion of the right coronary artery (RCA) or the left circumflex (LCX) coronary artery and the outcome depends on the culprit artery involved. The presence of significant ST elevation $>2\text{mm}$ in leads V5 and V6 with inferior wall AMI is a sensitive and specific sign of mega-artery-related infarction with expected high ischemic risk (Assali et al., 1998). Thus, the prediction of a culprit artery based on electrocardiography is of clinical importance. Our findings have important clinical implications to predict the infarct related artery (RCA or LCX) by comparing the degree of ST elevation in lead V6 with that of lead III in inferior wall STEMI.

Leads V5 and V6 face the posterolateral wall adjacent to the apex of the left ventricle (Cooksey et al., 1977). This region of the left ventricle is usually supplied by the larger obtuse marginal branch of the left circumflex artery (LCX), the atrioventricular branch of the right coronary artery, or the diagonal branch of the left anterior descending coronary artery. Therefore, the concurrent ST elevation in leads V5 and V6 during inferior AMI presenting with ST elevation in the inferior leads might reflect transmural ischemia extending to the posterolateral wall in addition to the inferior wall, which could be induced by either the right coronary artery (RCA) or the left circumflex artery (LCX) occlusion.

Fewer studies have suggested that ST elevation in leads V5 and V6 suggests the presence of the left circumflex artery (LCX) occlusion (Herz et al., 1997; Bairey et al., 1987; Assali et al., 1998; Kontos et al., 1997). In our study, among 62 patients with ST elevation in leads V5 and V6, the infarct-related artery was the right coronary artery in 41 patients (66%; $p=0.0001$) and left circumflex artery (LCX) in 19 patients (31%; $p=0.0001$). The ST elevation in leads III and V6 was strongly related to the infarct related artery. The ST elevation in lead III $> V6$ indicated right coronary artery (RCA) occlusion in 37 patients (82%) with the sensitivity and specificity of 90% and 63%, respectively. Whereas the ST elevation in lead III $\leq V6$ indicated left circumflex artery (LCX) occlusion in 12 patients (71%) with sensitivity and specificity of 63% and 90% respectively. These findings are consistent with the results of Kosuge M et al. (Kosuge et al., 2012). In the presence of ST elevation in leads V5 and V6 in inferior wall AMI, we believe that these simple ECG criteria are useful for predicting the infarct related artery (IRA) in the clinical practice to evaluate the high-risk. Lead III faces the right inferior wall, and is usually supplied by the right coronary artery distribution (Kosuge et al., 1998; Herz et al., 1997; Zimetbaum et al., 1998; Hasdai et al., 1995; Kontos et al., 1997; Zimetbaum and Josephson, 2003; Wagner et al., 2009). We, therefore,

selected lead III as an illustrative inferior lead in our study. ST segment elevation in lead III $> V6$ might suggest that the severity of transmural ischemia as evaluated by the degree of ST-segment elevation was comparatively greater in the right inferior wall to the posterolateral wall. This might suggest involvement of the mega-artery right coronary artery occlusion, irrespective of the site of occlusion, as shown in the present study. A similar findings were seen in Kosuge M. et al. (Kosuge et al., 2012). In contrast, ST elevation in lead III $\leq V6$ indicates the similar in the right inferior wall and the posterolateral wall or greater than in the latter. In the present study, 12 patients (71%) had LCX occlusion with ST elevation in lead III $\leq V6$. The ECG changes during LCX artery occlusion is highly variable because of its anatomical distribution. The LCX supplies blood to the posterior, posterolateral, posteroinferior, or posteroinferolateral myocardium (Dunn et al., 1984). During LCx occlusion, posterolateral wall involvement is associated with proximal occlusion because this region is supplied, at least in part, by the obtuse marginal branch of the LC artery (Hasdai et al., 1995; Dunn et al., 1984). When occlusion of the LCx artery is more distal, the posterolateral wall will not be involved, and the ST-segment vector will be oriented more inferior. Thus, the ST-segment will not be elevated in leads V5 and V6, and ST elevation in inferior leads could occur (Dunn et al., 1984). Thus, when assessing the greater risk area in patients with inferior wall AMI, the clinician should consider the presence of ST segment elevation in leads V5 and V6, which implies a large perfusion area of the left ventricle and the need for aggressive therapeutic strategies to improve myocardial reperfusion. ST-segment depression in the anterior leads V1-V3 in patients with inferior acute wall STEMI (ST elevation in II, III and aVF) have been shown to be useful for evaluating the infarct size and outcomes (Berger and Ryan, 1990; Evans et al., 1998; Peterson et al., 1996). In our study, the presence of ST-segment depression in leads V1-V3 with ST elevation in leads V5 and V6 with ST elevation in lead III $> V6$ and ST elevation in lead III $\leq V6$ was not of discriminatory value.

Tierala et al proposed that in inferior AMI, right coronary artery (RCA) might be the culprit artery in the ST elevation in lead III $> II$ and left circumflex artery (LCX) if the ST elevation in lead II $> III$ (Tierala et al., 2009). In our study, we reported that in patients with inferior AMI, the degree of ST elevation in the inferior leads, mainly leads III $> II$, was significantly greater in right coronary artery (RCA) occlusion than in left circumflex artery occlusion (81% vs 16%). Importantly, ST elevation in lead III $> II$ frequently supported ST elevation in leads V5 and V6 with ST elevation in lead III $> V6$ than the ST elevation in leads V5 and V6 with ST elevation in lead III $\leq V6$ (n=38; 84% vs n=9; 53%; $p=0.027$). However, ST elevation in lead II $> III$ favored ST elevation in leads V5 and V6 with ST elevation in lead III $\leq V6$ than the ST elevation in lead

III>V6 (n= 8; 47% vs n=7; 15%; $p=0.027$).

The present study had several limitations that need to be considered when interpreting the clinical implications of our findings. This was a small sample size, with retrospective study in a single center. The study didn't deal in predicting the lesion proximal or distal coronary artery occlusion and involvement of the mega artery. The study lacked the follow-up and complications like infection, postinfarction, angina, arrhythmias and mortality. The electrocardiographic findings in our study might differ considerably from those patients with previous episodes of myocardial infarction, late presentation. Thus, our findings cannot be generalized to all patients with inferior wall AMI and large scale studies are needed to confirm the present findings.

CONCLUSION

In conclusion, ST segment elevation in leads V5 and V6 suggests a greater risk area in patient with acute inferior wall myocardial infarction. The right coronary artery occlusion was most common in ST elevation in leads V5 and V6 with ST elevation in lead III>V6 and the left circumflex artery occlusion in ST elevation in leads V5 and V6 with ST elevation in lead III≤V6. And, while evaluating risk in an acute inferior wall myocardial infarction, we should also look for ST elevation in leads V5 and V6, as it signify a larger perfusion territory and demands for more aggressive reperfusion therapy.

REFERENCES

- Assali AR, Herz I, Vaturi M, Alder Y, Solodky A, Birnbaum Y, Sclarovsky S (1999). Electrocardiographic criteria for predicting the culprit artery in inferior wall acute myocardial infarction. *Am. J. Cardiol.* **84**:87-89.
- Assali AR, Sclarovsky S, Herz I, Adler Y, Porter A, Solodky A, Strasberg B (1998). Comparison of patients with inferior wall acute myocardial infarction with versus without ST-segment elevation in leads V5 and V6. *Am. J. Cardiol.* **81**:81– 83.
- Bairey CN, Shah PK, Lew AS, Hulse S (1987). Electrocardiographic differentiation of occlusion of the left circumflex versus the right coronary artery as a cause of inferior acute myocardial infarction. *Am. J. Cardiol.* **60**:456-459.
- Berger PB, Ryan TJ (1990). Inferior myocardial infarction: high-risk subgroups. *Circulation.* **81**:401– 411
- Braat SH, de Zwaan C, Brugada P, Coenegracht JM, Wellens HJ (1984). Right ventricular involvement with acute inferior wall myocardial infarction identifies high risk of developing atrioventricular nodal conduction disturbances. *Am. Heart J.* **107**:1183–1187.
- Chia BL, Yip JW, Tan HC, Lim TY (2000). Usefulness of ST elevation II/III ratio and ST deviation in lead I for identifying the culprit artery in inferior wall acute myocardial infarction. *Am. J. Cardiol.* **86**:341-343.
- Cooksey JD, Dunn M, Massie E (1977). Inferoposterior myocardial infarction. In: Cooksey JD, Dunn M, Massie E, eds. *Clinical Vactrocardiography and Electorocardiography*, 2. Chicago: Year Book Medical Publishers. pp. 391– 427.
- Dunn RF, Newman HN, Bernstein L, Harris PJ, Roubin GS, Morris J, Kelly DT (1984). The clinical features of isolated left circumflex coronary artery disease. *Circulation.* **69**:477– 484.
- Evans MA, Clements IP, Christian TF, Gibbons RJ (1998). Association between anterior ST depression and increased myocardial salvage following reperfusion therapy in patients with inferior myocardial infarction. *Am. J. Med.* **104**:5–11
- Fiol M, Cygankiewicz I, Carrillo A, Bayes-Genis A, Santoyo O, Gomez A, Bethencourt A (2004). Bayes de Luna A. Value of electrocardiographic algorithm based on "ups and downs" of ST in assessment of a culprit artery in evolving inferior wall acute myocardial infarction. *Am. J. Cardiol.* **94**:709-714.
- Hasdai D, Birnbaum Y, Herz I, Sclarovsky S, Mazur A, Solodky A (1995). ST segment depression in lateral limb leads in inferior wall acute myocardial infarction. *Eur. Heart J.* **16**:1549-1553.
- Herz I, Assali AR, Adler T, Solodky A, Sclarovsky S (1997). New electrocardiographic criteria for predicting either the right or left circumflex artery as the culprit coronary artery in inferior wall acute myocardial infarction. *Am. J. Cardiol.* **80**:1343-1345.
- Kontos MC, Desai PV, Jesse RL, Ornato JP (1997). Usefulness of the admission electrocardiogram for identifying the infarct-related artery in inferior wall acute myocardial infarction. *Am. J. Cardiol.* **79**:182– 184.
- Kosuge M, Ebina T, Hibi K, Iwahashi N, Morita S, Endo M, Maejima N, Gouhara M, Nagashima Z, Umemura S, Kimura K (2012). Implications of ST-Segment elevation in leads V5 and V6 in patients with reperfused inferior wall acute myocardial infarction. *Am. J. Cardiol.* **109**(3):314-319.
- Kosuge M, Kimura K, Ishikawa T, Hongo Y, Mochida Y, Sugiyama M, Tochikubo O (1998). New electrocardiographic criteria for predicting the site of coronary artery occlusion in inferior wall acute myocardial infarction. *Am. J. Cardiol.* **82**:1318-1322.
- Peterson ED, Hathaway WR, Zabel KM, Pieper KS, Granger CB, Wagner GS, Topol EJ, Bates ER, Simoons ML, Califf RM (1996). Prognostic significance of precordial ST segment depression during inferior myocardial infarction in the thrombolytic era: results in 16,521 patients. *J. Am. Coll. Cardiol.* **28**:305–312.
- Tierala I, Nikus KC, Sclarovsky S, Syvanne M, Eskola M (2009). Predicting the culprit artery in acute ST-elevation myocardial infarction and introducing a new algorithm to predict infarct-related artery in inferior ST-elevation myocardial infarction: correlation with coronary anatomy in the HAAMU Trial. *J. Electrocardiol.* **42**:120-127.
- Wagner GS, Macfarlane P, Wellens H, Josephson M, Gorgels A, Mirvis DM, Pahlm O, Surawicz B, Kligfield P, Childers R, Gettes LS, Bailey JJ, Deal BJ, Gorgels A, Hancock EW, Kors JA, Mason JW, Okin P, Rautaharju PM, van Herpen G (2009). American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology, American College of Cardiology Foundation, Heart Rhythm Society. AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram: part VI: Acute ischemia/infarction: a scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society. *J. Am. Coll. Cardiol.* **53**:1003–1011.
- Zimetbaum PJ, Josephson ME (2003). Use of the electrocardiogram in acute myocardial infarction. *N. Engl. J. Med.* **348**:933–940.
- Zimetbaum PJ, Krishnan S, Gold A, Carrozza II JP, Josephson ME (1998). Usefulness of ST-segment elevation in lead III exceeding that of lead II for identifying the location of the totally occluded coronary artery in inferior wall myocardial infarction. *Am. J. Cardiol.* **81**:918-919.