

Full Length Research Paper

Scrotal, testicular and semen characteristics of the mountain gazelle (*Gazella gazelle*) males

Al-Eissa Mohammed Saad^{1*}, Alhamidi A. R.², Mohamed Abd El Kader Sandouka³, Ibraheem A. S.¹, Hamad Al-Yahya⁴, AL-Farraj S.⁴, Saad Alkahtani⁴ and Al-Dahmash B.⁵

¹Department of Biology, College of Science, Hail University, Hail, Saudi Arabia.

²Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia.

³Department of Biology, Teachers College, King Saud University, Riyadh, Saudi Arabia.

⁴Department of Medical Laboratory, College of Health Sciences, King Saud University, Riyadh, Saudi Arabia.

Accepted 11 November, 2016

This study aimed to evaluate seasonal differences in the characteristics of various male reproductive organs and in semen quality. Ten adult males of mountain gazelles (*Gazella gazelle*) were used in this study during summer and winter. The results showed significant differences in July to September versus October to December for volume ($P \leq 0.05$), motility and viability ($P \leq 0.01$), progressive motility, and morphologic alteration ($P \leq 0.001$). Hence it was concluded that sperm characteristics are strongly influenced by season. This study therefore suggested that the suitable period for semen collection is the breeding season (October to December), which has a good quality for manipulation and long-term preservation of *G. gazelle* in Saudi Arabia.

Key words: *Gazelle*, reproduction, semen, sperm, cryopreservation.

INTRODUCTION

The Arabian sand gazelles (*Gazella subgutturosa marica*) is classified as vulnerable (VU) on the IUCN Red List (IUCN Red, 2009) and listed under appendix II of the convention on migratory species (CMS, 2005). One subspecies is listed on the IUCN Red List: the Arabian sand gazelle (*G. s. marica*) is classified as vulnerable (IUCN Red, 2009). Nevertheless, the following sub-species are all considered vulnerable: Hillier's goitered gazelles or Mongolian gazelles (*G. s. hillieriana*), Yarkand or Xingjian goitered gazelles (*G. s. yarkandensis*) and the Persian gazelles (*G. s. subgutturosa*) (East, 1992; Groombridge, 1993). Scientific references on the Arabian Sand Gazelles concern only the ethological aspects, reintroduction and some researches on veterinary and diseases (KKWRC, 2008). No data has been published on reproductive physiology and sperm characteristics

accept that was by (Al-Eissa, 1997, 2007a, 2007b).

Moreover, some studies described the reproductive season of Arabian sand gazelles and compared its reproductive cycles to Persian gazelle (*Gazella subgutturosa subgutturosa*) (Sempere, 2001). However, the length of gestation for Arabian sand gazelles is about 148 to 160 days compared to gestation length between 160 to 170 days for *G. cuvieri*, *G. dorcas*, and *G. leptoceros*, as mentioned by Al-Eissa (in press). In addition, evaluation of spermatozoa is an important factor that must be accurately analyzed to ensure the use of animals with good fertility. These evaluations often include measuring volume, concentration, motility, morphology, and other characteristics that influence sperm functionality.

Since it is important to improve the knowledge on reproductive male characteristics for protection of this species from extinction, this study was undertaken to investigate some testicular and sperm characteristics of the mountain gazelles and to find out which are the suitable months for semen quality collection, manipulation

*Corresponding author. E-mail: dr.aleissa@gmail.com. Tel: +699505782279.

and long-term preservation.

MATERIALS AND METHODS

Experimental animals

Ten adult males of mountain gazelles (*Gazella gazelle*) were used in this study. Animals were about three years old and around 14 kg weights. Animals were maintained and managed by the KKWRC, which is located in Riyadh, Saudi Arabia. Animals were transferred to an enclosure of 150 ha, while some animals were kept in small pens (0.5 ha) for research purposes. All animals were fed on dry alfalfa (*Medicago sativa*; min. 15% crude protein) and 1.5 L of water; quantities 10 to 15% above their average daily requirements (Ostrowski, 2006).

Extender

Various cryopreservation media (diluent) and Triladyl, a commercially available extender (Diluent A; Mnitub, Tiefenbach, Germany), were used.

Semen collection and processing

The animals were captured by using trap made with canvas around food point, then were caught manually inside the trap. The animal's heads were covered immediately on capture, and no stress was noted during handling. This study was conducted during 2008 in two different periods: July to September and October to December. Semen was collected from the males using the electroejaculation procedure described by Al-Eissa (2007a, 2007b), which typically involved less than 10 stimulations up to maxima of 4.5 to 7.0 volts as described by Holt et al. (1988, 1996). The animals were anaesthetized using xylazine (8.4 ± 1.5 mg/kg body weight) and ketamine hydrochloride (6.9 ± 1.5 mg/kg body weight). Testicular measurements were taken with a caliper and a tape after restraining the gazelle with a single intravenous administration of xylazine (8.4 ± 1.5 mg/kg body weight) and ketamine hydrochloride (6.9 ± 1.5 mg/kg body weight).

Testicular length was measured by placing the fixed arm of the caliper at the proximal end and the sliding arm at the distal end of the testis. Care was taken to exclude the epididymides. Thickness or depth was measured by placing the fixed arm of the caliper at the anterior aspect, and the sliding arm at the posterior aspect of each testis, at the point of maximum depth. Width of each testis was measured by sliding the other testis up in the scrotum and placing one arm of the caliper at the medial aspect and other at the lateral aspect, at the point of maximum width. For measurement of scrotal circumference the testicles were pushed firmly into the bottom of the scrotum by placing the thumb and fingers laterally on the side of neck of the scrotum and pushing ventrally. A flexible cloth tape was formed into a loop and slipped over the scrotum, and scrotal circumference was measured in centimeters by pulling the tape snugly around its greatest diameter (Elmore et al., 1976). Testes volume was calculated by using the formula for volume of an ellipsoid as described by Pant et al. (2003). Weight of the testes was calculated by multiplying volume with 1.038, which is the approximate density of testicular tissue (Amann, 1990). The seminal collection was obtained by electroejaculation as mentioned earlier. Volume was measured in a sterile glass tube graduated with 0.1 ml optical visible intervals, then, semen was diluted 1:100 with Triladyl and placed in water bath at 37°C. The evaluation was performed at the laboratory within 1 h.

Semen evaluation

Sperm volume, concentration and subjective scores of motility (wave motion, individual and progressive motility) were determined within 30 to 60 min of collection. The volume was measured by aspirating semen with a micropipette. Concentration was estimated using a haemocytometer. Semen was diluted (1:200) with an eosin solution (Madan et al., 1994). Motility was assessed by placing 10 µl of sperm suspension between a glass slide and a 22 mm × 22 mm cover slip, pre-warmed to 37°C. The proportion of motile spermatozoa and of spermatozoa with progressive motility (among motile ones) in each sample was assessed subjectively. In addition, motility was rated by using a scale 0 to 5 (0: non-motile; 5: highly motile), while the individual and progressive motility were determined by using eosin solution by placing 10 µl of sperm suspension on the slide without cover slip.

Sperm plasma membrane integrity was referred to viability by using acridine orange (AO), ethidium bromide (EB) and FluroQuench. Fluorescent dye has been used for analysis of DNA integrity of somatic cell nuclei and mammalian sperm nuclei. The AO staining ability of mammalian sperm was investigated to clarify the differences of its fluorescent emissions according to Kosower et al. (1992). However, stainability of sperm nuclei depends on the number of S-S bonds in sperm nuclear chromatin, and nuclear maturation of sperm can be analyzed by using AO fluorescence dye. A semen sample was mixed with 5 L of AO and incubated at 37°C for 10 min, then the sperms were evaluated with a n epifluorescence microscope (Nikon, Eclipse E 600 w.), at 1000 × magnification. Green spermatozoa (AO) were considered as alive, while those partially or completely red (EB) were considered dead. The percentage of plasma membrane intact spermatozoa was determined on at least 200 spermatozoa. Percentage of morphologic alterations was determined by analysis of sperm with normal morphology by using Spermac stain Enterprises, Wellington, (South Africa), (Baran et al., 2004; Schafer and Holzmann, 2000). A total of 200 cells were investigated by using the phase contrast microscopy (1000×).

Statistical analysis

Scrotal circumference and testicular diameters were statically analyzed by using *t*-test. Volume, concentration, motility, progressive motility, viability and morphologic alterations were analyzed by using one-way ANOVA test. Correlations were done for these parameters using SPSS version 16 (Sendecor and Cachran, 1980).

RESULTS

Testicular size as a proportion of body weight varies widely. In the mammalian of Saudi Arabia, there is a tendency for testicular size to correspond with multiple mates. Production of testicular output sperm and spermatogenic fluid is also larger in polygamous animals, possibly a spermatogenic competition for survival. Also, testis weight varies in seasonal breeders. In this study, the testicular measurement was summarized in Table 1.

Sperm characteristics, viability and sperm morphologic alterations during July to September and October to December periods are reported in Table 2. The statistical analysis showed significant differences in July to September versus October to December for volume ($P \leq 0.05$), motility and viability ($P \leq 0.01$), progressive

Table 1. Scrotal and testicular measurements in July to September period and October to December period, considered as non-breeding and breeding season, respectively.

| Measurement | July to September means (\pm S.D.) | October to December means (\pm S.D.) |
|--------------------------------------|---------------------------------------|-----------------------------------------|
| Scrotal circumference(cm) | 10.4 \pm 0.14 | 12.4 \pm 0.12 |
| Testis length (cm), right testicle | 3.8 \pm 0.28 | 4.0 \pm 0.24 |
| Testis length (cm), left testicle | 3.3 \pm 0.31 | 3.7 \pm 0.21 |
| Testis width (cm) , right testicle | 1.8 \pm 0.21 | 2.2 \pm 0.31 |
| Testis width (cm), left testicle | 1.5 \pm 0.41 | 1.9 \pm 0.11 |
| Testicular volume (cm ³) | 87 \pm 2.1 | 103.8 \pm 1.7 |

Table 2. Seminal characteristics calculated at 1, 10 and 24 h July to September period and October to December period considered as non-breeding and breeding season, respectively.

| Parameters | July to September means (\pm S.D.) | | | October to December means (\pm S.D.) | | |
|-----------------------------------------------|---------------------------------------|----------------|---------------|-----------------------------------------|----------------|----------------|
| | 1 | 10 | 24 | 1 | 10 | 24 |
| Time post-collection (h) | | | | | | |
| Volume (ml) | 0.19 \pm 0.026 | - | - | 0.418 \pm 0.077 | - | - |
| Concentration (\times 10 ⁶ /ml) | 574.17 \pm 108.2 | - | - | 910 \pm 59.89 | - | - |
| Total motility (%) | 49 \pm 10.46 | 20 \pm 9.5 | 0 | 75.33 \pm 4.82 | 32.1 \pm 6.4 | 53.7 \pm 3.5 |
| Progressive motility (%) | 37 \pm 8.16 | 5.0 \pm 2.1 | 0 | 73 \pm 4.53 | 30.2 \pm 2.1 | 43.9 \pm 2.9 |
| Viability (%) | 48.3 \pm 9.03 | 18.8 \pm 6.1 | 4.7 \pm 2.5 | 85.33 \pm 3.3 | 38 \pm 4.2 | 10.9 \pm 4.5 |

motility, and morphologic alteration ($P \leq 0.001$). Testicular diameters in July to September and October to December periods were determined and a light asymmetry between right and left testis in every single animal was observed. The scrotal circumference was significantly different in July to September than in October to December periods. More also, semen color was watery/opalescent during July September and creamy during October to December. Correlation coefficients of semen characteristics with testicular measurements and live weight are shown in Table 3. Scrotal circumference and width showed the largest coefficients ($r = 0.8$ to 0.85) with the semen volume and the total sperm number in the ejaculate, followed by the number of live and normal sperm in semen ($r = 0.70$). Correlation coefficients of these attributes with the testicular length were lower ($r = 0.45$ to 0.50) and barely reached the 5% significant level.

DISCUSSION

The limitations of this research were the scant number of valued subjects (Chemineau, 1986), the large extension of the protected area and the restricted number of captures (Delgadillo et al., 1991) in comparison with the attempts. According to Karagiannidis et al. (2000), a significant relationship between testicular diameters, ejaculate volume and breeding season exists in male. This correlation indicated that the animals in reproductive activity with repeated coupling showed a reduction in

semen volume for ejaculate. Other semen parameters values reported in bibliography in small ruminants were: volume 536 to 3786 μ L, concentration 358 to 922 $\times 10^6$ spermatozoa/ml, motility 53 to 81% and sperm abnormalities percentage 19.4 to 30% (Howard et al., 1983; Cassinello et al., 1998; Garde et al., 2003).

In this study, the mountain gazelles (*G. gazella*) semen characteristics were comparable. In fact, semen concentrations in mountain gazelle were 574 \pm 108 and 910 \pm 59 $\times 10^6$ spermatozoa/ml, respectively, in July to September and October to December periods. Seminal quality parameters in Mountain gazelle during October–December period showed a TM of 49 \pm 10.46% and PM of 37 \pm 8.16% after 1 h maintenance in Equitainer which was preserved at 54% after 34 h. In July to September period, the values showed a poor quality of semen, but the decrease was comparable with other studies. Value of progressive motility in October to December (73 \pm 4.53) was significantly greater than in July to September period (37 \pm 8.16) because some studies reported that influence of motility was strongly correlated with IVF rates on the outcome of *in vitro* fertilization/ intracytoplasmic sperm injection with fresh vs. frozen testicular sperm (Park et al., 2003). So it could be suitable a semen collection for long-term semen preservation in the months of October to December. The degree of fluctuating asymmetry is positively related to the coefficient of inbreeding and negatively related to the proportion of normal sperm, suggesting that it is a reliable indicator of genetic stress and of ejaculate quality (Roldan et al., 1998).

Table 3. Correlation coefficients of semen characteristics with testicular measurements and live weight in the mountain gazelle.

| Parameters | Scrotal circumference | Testis width | Testis length | Live weight |
|---------------------------------|-----------------------|--------------|---------------|-------------|
| Semen volume | 0.85*** | 0.79** | 0.51 | 0.73** |
| Sperm concentration | 0.32 | 0.36 | 0.54* | 0.35 |
| Percent live sperm | 0.13 | 0.10 | -0.04 | 0.01 |
| Percent abnormal sperm | -0.16 | -0.16 | -0.15 | 0.01 |
| Total sperm number | 0.86** | 0.83*** | 0.63* | 0.76** |
| Number of live sperm | 0.71** | 0.66** | 0.45 | 0.61* |
| Number of live and normal sperm | 0.75*** | 0.72** | 0.52 | 0.63* |

*Significant at $P < 0.05$; **significant at $P < 0.01$; ***significant at $P < 0.001$.

In mountain gazelle there seems to be a clear seasonal influence on both semen production and characteristics as it is also reported in some antelope breeds (Asher et al., 1993). In other antelope like goitered gazelles (*G. subgutturosa subgutturosa*) living in an arid environment conditions, intense sexual activity was observed during the breeding season (November to December), male gazelles control their harem, moving around with it when population density is low, but when population density is high some males become territorial and participate in reproduction, while others remain mobile in non-reproductive bachelor groups (Mambetdjumaev, 1970; Gorelov, 1972; Djenvorov, 1984; Blank, 1985). During the summer, most activity takes place in the late afternoon and early morning, consisting of leisurely walking and simultaneous grazing. At midday, herds take shelter in the shade, where they excavate shallow oval-shaped pits to lie in. During the cooler winter months, this midday break is significantly reduced, and sometimes even eliminated. If disturbed from its shelter, a goitered gazelle rapidly flees for 200-300 meters, pausing to assess the danger from this distance. A broad circular path is then taken back to the original resting spot. Extremely speedy, these gazelles can run up to 60 kmph/36 mph. Each animal generally consumes about 30% of its body weight in green matter per day, and can derive most of its needed moisture from it. In the spring and summer, groups may travel to water sources, but even still they rarely drink daily. Herds cover 10-30 kilometers per day in the winter, with these distances being reduced nearly tenfold in summer. Throughout much of their range, mountain gazelle undergo seasonal migrations. During the breeding season, adult males become territorial, using dung middens placed at strategic locations to indicate ownership. At this time, males emit hoarse bellows, and glandular activity increases significantly, with the result that the male is often seen smearing secretions on objects (Heptner et al., 1989) and (Walther, 1990). In contrary, (Al-Eissa et al., 2007a) reported a breeding season in late summer and autumn with high quantity (volume and concentration) and quality (percentage of motile and progressive spermatozoa, percentage of abnormal spermatozoa).

In the males, the breeding season in temperate areas seems to be different from the mountain gazelle. A possible hypothesis could be related to the Mountain gazelle different sensitivity to photoperiod or other influences on melatonin release. The studied periods (July to September and October to December) represented non-breeding and breeding season respectively, if compared with other study as carried out by Al-Eissa, (2007a), the results were conformable with the analysis of reproductive characteristics during whole year (Al-Eissa, 2007a). Semen characteristics of mountain gazelle are highly influenced by season. The acceptable semen production to try cryopreservation programs was produced in October to December, but more investigations are needed in order to identify the exact breeding period. Frozen semen could be utilized in artificial insemination for genetic studies, so as to increase genotypic variability and resolve the problem of rupicapra species extinction.

Conclusion

Since sperm characteristics are strongly influenced by season, this study therefore suggests that the suitable period for semen collection is the breeding season (October to December), which has a good quality for manipulation and long-term preservation of *G. gazelle* in Saudi Arabia.

ACKNOWLEDGEMENT

The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for funding the work through the research group project no. RGP-VPP-018.

REFERENCES

- Al-Eissa MS, Alhomida AS (1997). "A study of the distribution of total, free, short-chain acyl and long-chain acyl carnitine in whole blood and plasma of Arabian Sand Gazelle (*Gazelle Subgutturosa marica*)", *Comparat. Haematol. Int.*, 1: 65-69.

- Al-Eissa MS (2007a). Assessment of reproductive efficiency of the Arabian Sand Gazelle males (*Gazella subgutturosa marica*) and semen cryopreservation". PhD Thesis, King Saud University, College of Science, Department of Zoology. pp. 25-36
- Al-Eissa MS, Alhamidi AR, Khandil S (2007b). Semen cryopreservation using triadyl and tris diluents of the Arabian Sand Gazelle males (*Gazelle Subgutturosa marica*) Arabian Gulf J. Sci. Res., (4)25: 199-206.
- Al-Eissa MS, Mohamed Abd El Kader S, Saad H, Ashraf AQ (2009). Reproductive characteristics of The Arabian Sand Gazelles (*Gazella subgutturosa marica*). J. Audi Society Agric. Sci., (9): 70-80.
- Amann RP (1990). Management of the bull to maximize sperm output. Proceedings of the 13th Technical conference on AI Reproduction. pp.
- Asher GW, Fisher MW, Fennessy PF, Mackintosh CG, Jabbour HN, Morrow CJ (1993). Oestrous synchronization, semen collection and artificial insemination of farmed red deer (*Cervus elaphus*) and fallow deer (*Dama dama*). Anim. Reprod. Sci., 33: 241-265.
- Baran AI, Evrim BS, Evecen M, Demir K, Kamuran (2004). Use of spermac staining technique in the determination of acrosomal defects in cat semen. Turk. J. Vet. Anim. Sci., 28: 519-525.
- Blank DA (1985). Specificity of social and reproductive behaviour of gazelles (*Gazella subgutturosa*) in the valley of Ili river. Zoologicheskii J., 64: 1059-1070.
- Cassinello J, Abaigar T, Gomendio M, Roldan ERS (1998). Characteristics of the semen of three endangered species of gazelles (*Gazelle dama mhorr*, *G. dorcas neglecta* and *G. cuvieri*). J. Reprod. Fertil., 113: 35-45.
- CMS (2005). Convention on Migratory Species. <http://www.cms.int>
- Chemineau P (1986). Sexual behaviour and gonadal activity during the year in the tropical Creole meat goat. II. Male mating behaviour, testis diameter, ejaculate characteristics and fertility. Reprod. Nutr. Dev., 26: 453-460.
- Delgado JA, Leboeuf B, Chemineau P (1991). Decrease in seasonality of sexual behaviour and sperm production in bucks by exposure to short photoperiodic cycles. Theriogenology, 36: 755-770.
- Djevevov VV (1984). Goitred Gazelle of Barsakelmes Island. Alma-Ata: Nauka. pp. 144.
- East R (1992). Conservation status of Antelopes in Asia and the Middle East, part 1. Species, 19: 23 - 25.
- Elmore RG, Bierschwal CJ, Youngquist RS (1976). Scrotal circumference measurements in 764 beef bulls. Theriogenology, 6: 485-489.
- Garde J, Soler AJ, Cassinello J, Crespo C, Malo AF, Espeso G, Gomendio M, Roldan ERS (2003). Sperm Cryopreservation in three species of endangered gazelles (*Gazella cuvieri*, *G. dama mhorr*, and *G. dorcas neglecta*). Biol. Reprod., 69: 602-611.
- Gorelov JK (1972). Reproduction of goitred gazelles and the problem of its restoration in Badhis. Theriology, 1: 420- 424.
- Groombridge B (1993). 1994 IUCN red list of threatened animals. International Union for Conservation of Nature and National Resources, Gland, Switzerland and Cambridge, U.K.
- Heptner VG, Nasimovich AA, Bannikov AG (1989). Mammals of the Soviet Union. New York: E.J. Brill., pp. 101-124.
- Holt WV, Moore HD, North RD, Hartman TD, Hodges JK (1988). Hormonal and behavioural detection of oestrus in blackbuck, Antelope cervicapra, and successful artificial insemination with fresh and frozen semen. J. Reprod. Fertil., 8: 2 717-725.
- Holt WV, Abaigar T, Jabbour H (1996). Oestrus synchronization, semen preservation and artificial insemination in the Mohor gazelle (*Gazella dama mhori*) for the establishment of a genome resource bank programme. Reprod. Fertil. Dev., 8: 1215-1222.
- Howard JG, Wildt DE, Chakraborty PK, Bush M (1983). Reproductive traits including seasonal observations on semen quality and serum hormone concentrations in the dorcas gazelle. Theriogenology, 20: 221-234.
- IUCN Red List (2009). <http://www.iucnredlist.org>
- Karagiannidis A, Varsakely S, Karatzas G (2000). Characteristics and seasonal variations in the semen of Alpine, Saanen and Damascus goat bucks born and raised in Greece. Theriogenology, 53: 1285-1293.
- KKWRC (2008). National Commission for wildlife Conservation and Development (NCWCD). Riyadh, Saudi Arabia.
- Kosower NS, Katayose H, Yanagimuchi RJN (1992). Thiol disulfide status and acridine orange fluorescence of mammalian sperm nuclei. J. Androl., 13: 342-348.
- Madan HC, Tamuli, Watson P (1994). Use of simple staining technique to distinguish acrosomal changes in the live sperm sub-population. Anim. Reprod. Sci., 35: 247-254.
- Mambetdumaev A (1970). *Goitred Gazelles*. Tashkent, Fan, pp. 196.
- Ostrowski S, Williams JB, Me'sochina P, Sauerw H (2006). Physiological acclimation of a desert antelope, Arabian oryx (*Oryx leucoryx*), to long-term food and water restriction. J. Comp. Physiol., B 176: 191-201.
- Pant HC, Sharma RK, Patel SH, Shukla HR, Mittal AK, Kasiraj R, Misra AK, Prabhakar JH (2003). Testicular development and its relationship to semen production in Murrah buffalo bulls. Theriogenology, 60: 27-34.
- Park YS, Lee SH, Song SJ, Jun JH, Koong MK, Seo JT (2003). Influence of motility on the outcome of in vitro fertilization / intracytoplasmic sperm injection with fresh vs. frozen testicular sperm from men with obstructive azoospermia. Fertil. Steril. 80: 526-530.
- Roldan ERS, Cassinello T, Abaigar, Gomendio M (1998). Inbreeding, fluctuating asymmetry, and ejaculate quality in an endangered ungulate. Proc. R. Soc. Lond. B., 265: 243-248.
- Schafer S, Holzmann A (2000). The use of transmigrating and Spermac stain to evaluate epididymal cat spermatozoa. Anim. Reprod. Sci., 59: 201-211.
- Sempere J, Brown N, Pereladova OB, Bahloul K, Lacroix A, Soldatova N (2001). Comparative analysis of reproductive cycles in female Persian gazelle (*Gazella subgutturosa subgutturosa*) (Central Asia) and Sand Gazelle (*Gazella subgutturosa marica*) (Arabian Peninsula). Gen. Comp. Endocrinol., 121: 57-65.
- Sendecor GW, Cochran WGH (1980). Statistical Methods 7thed, The Iowa State Univ. Press, Ames, Iowa, pp. 507.
- Walther FR (1990). Gazelles and related species. In Grzimek's Encyclopedia of Mammals. Edited by S. P. Parker. New York: McGraw-Hill., 5: 462-484.