

Full Length Research Paper

Evaluation of the efficacy of endophytic *Bacillus amyloliquefaciens* against *Botryosphaeria dothidea* and other phytopathogenic microorganisms

Xiang-Tian Yin¹, Li-Na Xu¹, Liang Xu¹, Su-Su Fan¹, Zhen-Yu Liu^{1*} and Xing-Yao Zhang²

¹College of Plant Protection, Shandong Agricultural University, 271018, Tai'an, Shandong Province, China. ²Research Institute of Forest Ecology Environment and Protection, Chinese Academy of Forestry, 100010, Beijing, China.

Accepted 13 January, 2019

Endophytic bacteria were isolated from poplar trees in Shandong province, China, where canker disease caused by *Botryosphaeria dothidea* has reached epidemic proportions. On the basis of the results of classical bacteriological tests and sequence analysis of the 16S rRNA gene, the PEBA20 strain isolated from poplar was identified as a strain of *Bacillus amyloliquefaciens*. *In vitro* PEBA20 inhibited the mycelium growth of *B. dothidea* and canker disease incidence on the cut shoots of poplar 60% than that in the control 100%. The optimum culture conditions for the propagation of the antagonistic strain against *B. dothidea* were determined using an orthogonal experiment design. The strain showed antifungal and antibacterial activity against a number of microorganisms, including mycelial fungi, yeast, gram-positive and gram-negative bacteria. *TasA* and *aiiA*—2 genes associated with antagonistic activity were detected in PEBA20. The results of our study indicate that *B. amyloliquefaciens* PEBA20 has the potential to serve as a biological control agent for the poplar canker disease caused by *B. dothidea* and for diseases caused by other phytopathogens.

Key words: *Bacillus amyloliquefaciens*, endophytic bacteria, isolation, biological control agent.

INTRODUCTION

Poplar (*Populus* spp.) is a widely cultivated plant and is used as a source of wood and as an energy crop for the production of renewable sources of energy. Poplar canker disease is caused by *Botryosphaeria dothidea* (Moug.:Fr.) Ces. and De Not. and is an important fungal disease of the stem in China and other countries throughout the world (Morgan-Jones and White, 1987). Disease management strategies for poplar canker typically include the use of genetically resistant varieties of the plant, integration of select culture practices, application of chemical pesticides, and combinations thereof. However, the use of some of these practices is not always viable, and the continuous use of pesticides has caused severe ecological problems. Owing to their

environment-friendly nature, biological control methods have emerged as an alternative to chemical treatments for plant disease control (Daughtrey and Benson, 2005; Strange and Scott, 2005). Microorganisms can colonize the tissues of healthy plants. Such endophytic bacteria have been reported to prevent disease development by controlling the spread of plant pathogens or by enhancing plant resistance (Stein, 2005; Ryan et al., 2008). Therefore, there is a considerable potential for finding new and beneficial endophytic bacteria that can serve as biocontrol agents.

Few studies have reported the biological control of poplar canker disease. Previous studies have revealed some strains, including *Bacillus subtilis*, *Bacillus pumilus*, *Pseudomonas fluorescens*, *Trichoderma* spp., *Eudarluka quinquesepata*, and *Eudarluka biconia*, that are potential biocontrol agents against *B. dothidea* (Katamoto, 1986; Jin and Shen, 1989). However, these strains were isolated from either soil or the rhizosphere and

*Corresponding author. E-mail: cosmosliu@163.com. Tel: 86-538-8247781-8805. Fax: 86-538-8241324.

Table 1. Antagonistic activity of *Bacillus amyloliquefaciens* PEBA20 against mycelial fungi.

Indicator fungi	Inhibitory rate (%) ^a	
	Treated with bacterial suspension of PEBA20	Treated with fermentation filtrate of PEBA20
<i>B. dothidea</i>	69.95	73.2
<i>Fusarium oxysporum</i>	85.94	63.55
<i>Fusicladium tremulae</i>	45.91	40.99
<i>Gloeosporium fructigenum</i>	75.75	64.87
<i>Glomerella cingulata</i>	60.35	46.05
<i>Myrothecium roridum</i>	38.44	26.26
<i>Phytophthora capsici</i>	81.06	71.3
<i>Rhizoctonia solani</i>	39.05	19.2
<i>Sphaceloma euonymi-japonici</i>	55.46	52.05
<i>Trichothecium roseum</i>	64.02	53.7
<i>Valsa sordida</i>	89.37	82.62

^a Values are the mean of 4 replicates; the formula for inhibitory rate is as follows: Inhibitory rate (%) = [(colony radius in controls colony radius in treatments)/colony radius in controls] x 100.

Table 2. Antagonistic activity of *Bacillus amyloliquefaciens* PEBA20 against yeast and bacteria.

Indicator yeast and bacteria	Diameter of the inhibitory zone (mm) ^a	
	Treated with bacterial suspension of PEBA20 ^b	Treated with fermentation filtrate of PEBA20 ^c
<i>Saccharomyces cerevisiae</i>	28.51	9.04
<i>Xanthomonas campestris</i>	19.14	15.76
<i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>	9.97	9.21
<i>Ralstonia solanacearum</i>	13.30	13.31
<i>Clavibacter michiganense</i> subsp. <i>sepedonicum</i>	16.61	16.68
<i>Staphylococcus aureus</i>	10.01	5.1
<i>B. subtilis</i>	13.57	6.67
<i>B. cereus</i>	18.95	9.45
<i>B. megaterium</i>	18.52	19.05

^a Values are the mean of 4 replicates; the diameter of the inhibitory zone includes the diameter of the well (5 mm), ^b Treatment by adding 20 l of bacterial suspension in a well with diameter 5 mm, ^c Treatment by adding 20 l of fermentation filtrate in a well with diameter 5 mm.

phyllosphere; to date, no study has reported the isolation of endophytic antagonistic bacteria from poplar. Although endophytic bacteria from poplar plants have been reported, these studies did not assess the biological control mediated by these organisms (Doty et al., 2005; Frey et al., 2008; Ulrich et al., 2008). In this study, we isolated a bacterial strain from poplar as a potential biocontrol agent and assessed its inhibitory activity against *B. dothidea*.

MATERIALS AND METHODS

Bacterial and fungal strains

The fungi and bacteria used as indicators for evaluating antagonistic activity are listed in Tables 1 and 2. The fungi were stored at 4°C and the bacteria, at -20°C. Before being used for

evaluating antagonistic activity, the fungal cultures were routinely incubated on potato dextrose agar (PDA) at 25°C, while the bacterial cultures were incubated on nutrient agar (NA) at 28°C.

Isolation of endophytic bacteria and screening for antagonistic activity against *B. dothidea*

Endophytic bacteria were isolated from five 6-year-old poplar trees (*Populus tomentosa* Carr.). A tissue sample (10 mm in depth and 40 mm in length and breadth) comprising the epidermis, phloem, and xylem was cut from the outer surface of the tree trunk at a height of 1.5 m from the ground. The samples were surface sterilized with 70% alcohol and the epidermis and phloem were discarded. Then, the remaining tissue was surface sterilized with 70% alcohol for 1 min and 2% sodium hypochlorite for 10 min and rinsed 5 times in sterile distilled water. Hundred microliters of the last wash of each sample was spread onto NA plates to check the sterility of the sample. Then, the tissue samples were aseptically ground in sterile distilled water. The supernatant was diluted up to 0.01 g·ml⁻¹, pour-

plated on NA plates, and incubated at 28°C. After incubation, individual bacterial colonies were selected and subcultured on NA plates. An *in vitro* dual culture technique was used to screen the isolates for antagonistic activity against *B. dothidea* (Katamoto, 1986; Ryan et al., 2008). A mycelial plug of *B. dothidea* (diameter, 5 mm) and a paper disc (diameter, 5 mm) dipped in overnight cultures of the isolates were placed 25 mm apart on a Petri dish (90 mm) containing PDA. Inhibition of fungal growth was assessed on the basis of the presence or absence of an inhibition zone. Isolate 080120 showed strong inhibitory activities and was selected for further studies.

Bacterial characterization and 16S rRNA gene sequence analysis

Cultural, morphological, and physiological characteristics of isolate 080120 were assessed using conventional methods (Colwell and Grigorova, 1987; Holt, 1994). Genomic DNA was extracted by the cetyltrimethylammonium bromide (CTAB) method (Ausubel et al., 1992). For polymerase chain reaction (PCR) amplification of the 16S rRNA gene, primers P3—5'-AGAGTTTGATCMTGGCTCAG-3'—and P5—5'-TACGGYTACCTTGTACGACTT-3'—were designed using DNAMAN 6.0 on the basis of the previously described universal primers and the 16S rRNA gene sequences of *Bacillus* species in GenBank (Moreno et al., 2002). PCR amplification consisted of initial denaturation at 94°C for 5 min followed by 30 cycles of denaturation at 94°C for 60 s, annealing at 58°C for 60 s, elongation at 72°C for 90 s, and a final extension at 72°C for 10 min. Subsequently, the PCR product was subcloned and sequenced. Two clones from each independent PCR reaction were sequenced twice on both strands. Two independent PCR reactions were performed to obtain the consensus sequence. Nucleotide sequence alignment and analyses were performed using DNAMAN 6.0 and Molecular Evolutionary Genetic Analysis (MEGA) version 4.1.

Detection of *TasA* and *aiiA* genes associated with antagonistic activity

Detection of the *TasA* and *aiiA* genes was performed using PCR amplification (Lee et al., 2002; Branda et al., 2006). The PCR primers to detect these genes were designed using DNAMAN 6.0 on the basis of homologous sequences obtained from GenBank. The primers *aiiA*1—5'-GGAATTCATGACAGTAAARAARCT-3'—and *aiiA*2—5'-CCCTCGAGTATATAYTCHGGGAAC-3'—were used for *aiiA* amplification and *TasA*1—5'-GAATTCATGGGTATGAAAAAGAA-3'—and *TasA*2—5'-CTGCAGTTAATTTTATCCTCGCT-3'—were used for *TasA* amplification (*EcoR* site for *aiiA*1 and *TasA*1, *Xho* site for *aiiA*2 and *Pst* I site for *TasA*2 have been underlined). The *aiiA* gene was amplified under the following conditions: initial denaturation at 94°C for 5 min, 30 cycles of denaturation at 94°C for 60 s, annealing at 50°C for 45 s, elongation at 72°C for 60 s, and a final extension at 72°C for 7 min. For *TasA* amplification, touchdown PCR was performed with annealing at 56°C for 45 s after which the temperature was decreased to 52°C at the rate of 2°C per 2 cycles; this step was followed by 28 cycles at 50°C for 45 s. The amplified products were subcloned and sequenced. The sequence was compared with homologous sequences in GenBank.

Assessment of antifungal and antibacterial activity

To determine antifungal activity, a paper disc (diameter, 5 mm) dipped in overnight suspension of isolate 080120 was placed on a PDA plate at a distance of 50 mm from the mycelial plug (diameter, 5 mm) of the indicator fungi. For the fermentation filtrate test, the

bacteria were inoculated in 250 ml Erlenmeyer flasks containing nutrient broth (NB; 150 ml) and grown at 28°C and 180 rpm for 120 h. The cell-free fermentation filtrate thus obtained was previously added to the PDA plate at a concentration of 200 l of fermentation filtrate per 10 ml PDA in a Petri dish (90 x 15 mm). The mycelial plug of the indicator fungi was placed at the center of the PDA plate. Then, the PDA plates were incubated at 25°C for 3–5 days, depending on the growth rate of the fungi. The antagonistic activity against *Saccharomyces cerevisiae* and bacteria was assessed using well-diffusion inhibition assays (Cintas et al., 1995). The overnight cultures of indicator yeast and bacteria (100 l) were added to the base plates containing PDA and NA, respectively and incubated to obtain lawns of the organisms on the plates. A well (diameter, 5 mm) was bored into the base plate and overnight culture of isolate 080120 (20 l) or the cell-free fermentation filtrate was added to the well. The plates were then incubated at 28°C for 2 days. To establish the controls, the fungi and bacteria were incubated under the same conditions without the bacterial suspensions and cell-free fermentation filtrate. Four replications were performed for each treatment.

Optimization of culture conditions for the propagation of an antagonist strain

To determine the optimum culture medium for the propagation of an antagonist strain against *B. dothidea*, NB, Luria-Bertani (LB), BPY (beef, peptone, and yeast), and PD media were used. The inoculum of isolate 080120 was incubated in 250 ml Erlenmeyer flasks containing 150 ml of different culture media at 28°C and 180 rpm for 120 h. To determine the antifungal activity of the cell-free fermentation filtrate, the optimum medium was chosen for optimization of the culture conditions. The optimization of culture conditions for the propagation of the antagonist strain was performed using an orthogonal experimental design. Five factors at 5 levels, viz., volume of the medium (50, 100, 125, 150 and 175 ml), initial pH (5, 6, 7, 8 and 8.67), temperature (20.0, 24.2, 28.5, 32.8 and 37°C), shaking speed (100, 120, 140, 160, and 180 rpm) and cultivation time (24, 48, 72, 96 and 120 h) were selected and an orthogonal array layout of 5⁵ was performed. The antifungal activity of the cell-free fermentation filtrate was determined by the methods described above.

Cut shoots assay for suppression of poplar canker

The efficacy of isolate 080120 in suppressing canker caused by *B. dothidea* was assessed using stem cuttings of poplar. The basal end of healthy shoots of *P. tomentosa* (length, about 40 cm) were dipped in a bacterial suspension (10 CFU ml⁻¹) for 30 min. Then, the shoots were hydroponically cultured at room temperature (about 28°C) for a week. A mycelial plug of *B. dothidea* was inoculated on the stem and incubated at 28°C in a moisture chamber. Dark brown to black lesions were observed on the stem. This test was performed in June; 30 cuttings were used for the treatments and 10 were used as controls.

RESULTS

Isolation and identification of antagonistic bacteria

A total of 126 bacterial isolates were collected from the surface-sterilized internal tissues of poplar trees. The results of screening for antagonistic activity against




Figure 1. PCR amplification of *aiiA* gene from *B. amyloliquefaciens* PEBA20. Lane 1: products of PCR amplification, lane M: molecular weight marker.




Figure 2. PCR amplification of *TasA* gene from *B. amyloliquefaciens* PEBA20. Lane 1: products of PCR amplification, lane M: molecular weight marker.

obtained on NA medium; the colonies were circular, undulated, raised, rough, opaque, and greyish white in color with no diffusible pigment. Classical bacteriological tests showed that the isolate consisted of motile, aerobic Gram-positive long rods that showed endospore formation; were positive for catalase activity, oxidase activity, Voges-Proskauer test, gelatin hydrolysis, and starch degradation; and could utilize D-glucose, D-xylose, L-arabinose, and D-mannitol.

The partial 16S *rRNA* gene sequence was amplified by using the genomic DNA of isolate 080120; this sequence has been deposited in GenBank with accession number FJ685773. Basic Local Alignment Search Tool (BLAST) analysis and a neighbor-joining dendrogram constructed using MEGA revealed that the bacterium belonged to the genus *Bacillus* and was closely clustered together with *Bacillus amyloliquefaciens*. The amplified 16S *rRNA* gene sequence of isolate 080120 was most closely related to that of *B. amyloliquefaciens* FZB42 (GenBank accession number, CP000560) and showed 100% identity with the sequence from *B. amyloliquefaciens* FZB42. The sequence showed 99.87 to 99.93% identity with the sequences obtained from most of the *B. amyloliquefaciens* strains, whereas it showed 99.73 to 99.80% identity with the sequences obtained from most of the *B. subtilis* strains. On the basis of the results of the classical bacteriological tests and the analysis of the 16S *rRNA* gene, we concluded that the isolate was a strain of *B. amyloliquefaciens* and was named PEBA20.

Further, 2 genes—*aiiA* and *TasA*—from the genomic DNA of PEBA20 were amplified using PCR to facilitate gene sequencing to confirm the identity of the strain (Figures 1 and 2). The intact open reading frame (ORF) of the *aiiA* gene (GenBank accession number, FJ713589) consists of 753 base pairs, which encode a protein of 250 amino acid residues. The gene showed 87.0 to 99.3% identity with the homologues of the *aiiA* genes cloned from strains of *Bacillus thuringiensis*, *Bacillus cereus*, *Bacillus anthracis*, *Bacillus weihenstephanensis*, and *B. subtilis*. The ORF of the *TasA* gene (GenBank accession number, FJ713580) consists of 786 base pairs, which encode a protein of 261 amino acid residues. The gene showed 98.60% identity with the *TasA* gene of *B. amyloliquefaciens* FZB42 (GenBank accession number, CP000560) and 76.02% identity with 3 homologues from *B. subtilis* strains (GenBank accession numbers, BACJH642; AJ871386; and AL009126). The results of the gene sequence analysis indicated that PEBA20 has the *aiiA* and *TasA* genes and that it is a strain of *B. amyloliquefaciens*.

Antifungal and antibacterial spectrum

B. amyloliquefaciens PEBA20 showed antimicrobial activity against a wide range of fungi and bacteria. The mycelial fungi, yeast, and gram-positive and negative

B. dothidea showed inhibition zones with different areas for different isolates. Isolate 080120 showed strong inhibitory activity against *B. dothidea*. After incubation for 96 h, solitary bacterial colonies of isolate 080120 were




Figure 3. Normal hyphae of *B. dothidea* at the edge of a colony cultured on a potato dextrose agar (PDA) plate at 25°C for 120 h (magnification, 400 x).

bacteria tested in the study showed various degrees of sensitivity to bacterial suspensions or fermentation filtrates of PEBA20 (Tables 1 and 2). The efficacy of the bacterial suspension in inhibiting fungal growth *in vitro* ranged from 38.44 to 89.37%, while that of the fermentation filtrate ranged from 19.2 to 82.62%. The diameter of the inhibition zone after treatment with bacterial suspension and fermentation filtrate of PEBA20 ranged from 9.97 to 19.14 mm and from 5.1 mm (very weak effect) to 19.05 mm. However, no relationship was seen either among the different indicators or between the treatments using bacterial suspensions and fermentation filtrate.

Antagonistic activity against *B. dothidea*

Bacterial suspensions and the fermentation filtrate of PEBA20 inhibited the growth of *B. dothidea* *in vitro*, thereby showing the marked antagonistic activity of PEBA20 (Table 1). The inhibitory effect of the fermentation filtrate varied in different culture media; the maximum rate of inhibition (84.33%) was obtained when PEBA20 was incubated in NB, and the rate of inhibition decreased to 77.81, 75.2 and 60.31% when PEBA20 was incubated in LB, BPY, and PD, respectively. Therefore, NB medium was chosen as the optimum culture medium. The optimum culture conditions for the propagation of the strain with antagonistic activity against *B. dothidea* were as follows: 100 ml of NB in a 250 ml flask; initial pH, 5; temperature, 28.5°C; shaking speed, 140 rpm; and cultivation time, 120 h.

Aberrant hyphae were detected when the fungus were treated with either the bacterial suspensions or fermentation filtrate. Compared with the normal hyphae (Figure 3), the hyphae at the edge of the colony showed




Figure 4. Aberrant hyphae of *B. dothidea* treated with *Bacillus amyloliquefaciens* PEBA20 (magnification, 400×) * distorted hyphae at the edge of a colony cultured on potato dextrose agar (PDA) plate at 25°C for 120 h after treatment with bacterial suspensions of PEBA20 (a paper disc immersed in the bacterial suspension was placed on the PDA plate at a distance of 50 mm from the mycelial plug).

obvious hyphal aberration, such as increasing branches, swelling, and bulb formation in the middle and/or at the top (Figure 4). Our results indicated that the hyphal aberration increased with treatment duration; some hyphae had a bead-like appearance, while some formed cluster-like structures.

We used cut shoots of poplar to determine the efficacy of PEBA20 against *B. dothidea* infection. Our results showed that the incidence of poplar canker in the cut shoots treated with the bacterial suspension of PEBA20 (60%) was lower than that in the control (100%). In addition, the canker symptoms in the treated cut shoots were different from those in the controls. The diseased cut shoots treated with the PEBA20 bacterial suspension showed delayed incidence of typical canker symptoms and decrease in the average radius of the lesion.

DISCUSSION

Endophytic microorganisms have attracted the attention of researchers because of their potential to serve as biocontrol agents (Stein, 2005; Ryan et al., 2008). In this study, *B. amyloliquefaciens* strain PEBA20 was isolated from poplar. The efficacy of this strain as a biocontrol agent against poplar canker caused by *B. dothidea* was evaluated by *in vitro* bioassay and cut shoots assay. Our work provides evidence for the antagonistic activity of endophytic bacteria against *B. dothidea* infection in poplar (Katamoto, 1986; Doty et al., 2005; Frey et al., 2008; Ulrich, 2008). Since the PEBA20 strain was obtained from poplar, it has the advantage of being a “native” antagonist isolated from the same habitat in

which it is expected to be used for biological control.

Species of the genus *Bacillus*, particularly *B. amyloliquefaciens* and *B. subtilis*, have been shown to produce a range of antimicrobial dipeptides or cyclic lipopeptides. Some of the metabolites are strain-specific and may be associated with certain species and subspecies of *B. amyloliquefaciens* and *B. subtilis* (Stein, 2005; Romero et al., 2007). Although we did not detect the antibiotics produced by PEBA20 in the present study, the detection of antagonistic activity and aberrant hyphae indicated the production of antimicrobial compounds.

In this study, 11 mycelial fungi, 1 yeast, 5 Gram positive bacteria, and 3 Gram negative bacteria were found to be sensitive to PEBA20, thereby implying that PEBA20 has a wide spectrum of antimicrobial activity; however, since a large antimicrobial spectrum is an important attribute for a successful biocontrol agent, more fungi and bacteria should be tested in further studies to establish the antimicrobial spectrum of this strain. We also detected the *aiiA* and *TasA* genes in PEBA20. *AiiA*, encoded by the *aiiA* gene, is a quorum-quenching agent and can be a promising agent for biological control of bacterial diseases (Dong et al., 2001; Dong and Zhang, 2005). *TasA* is a spore-associated protein encoded by *TasA* in *B. subtilis* and is a protein with the property of antibacterial activity (Stover and Wong, 1999). In addition, *TasA* is a predominant protein component of the extracellular matrix of biofilms, which contribute to important biological function of bacteria (Branda et al., 2006). PEBA20 was found to possess *TasA* and *aiiA*, which suggested its potential antagonistic activity and indicated that it serves as a quorum-quenching agent in biological control of bacterial diseases. The results indicate that PEBA20 has immense potential to serve as a biocontrol agent.

ACKNOWLEDGEMENTS

The authors thank the anonymous reviewers for their valuable suggestions. This work was supported by grants from the National Natural Science Foundation of China (30972367) and the Advanced Postdoctoral Science Foundation of Shandong Province (200603059).

REFERENCES

- Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K (1992). Short protocols in molecular biology. 3rd edn. New York: John Wiley and Sons.
- Branda SS, Chu F, Kearns DB, Losick R, Kolter R (2006). A major protein component of the *Bacillus subtilis* biofilm matrix. *Mol. Microbiol.*, 59: 1229-1238.
- Cintas LM, Rodriguez JM, Fernandez MF, Sletten K, Nes IF, Hernandez PE, Holo H (1995). Isolation and characterization of pediocin L50, a new bacteriocin from *Pediococcus acidilactici* with a broad inhibitory spectrum. *Appl. Environ. Microbiol.*, 61: 2643-2648.
- Colwell R, Grigorova R (1987). *Methods in microbiology, current methods for classification and identification of microorganisms*. Orlando FL: Academic Press.
- Daughtrey ML, Benson DM (2005). Principles of plant health management for ornamental plants. *Annu. Rev. Phytopathol.*, 43: 141-149.
- Dong Y, Zhang L (2005). Quorum sensing and quorum-quenching enzymes. *J. Microbiol.*, 43: 101-109.
- Dong YH, Wang LH, Xu JL, Zhang HB, Zhang XF, Zhang LH (2001). Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature*, 411: 813-817.
- Doty SL, Doshier MR, Singleton GL, Moore AL, Van Aken B, Stettler RF, Strand SE, Gordon MP (2005). Identification of an endophytic *Rhizobium* in stems of *Populus*. *Symbiosis*, 39: 27-35.
- Frey B, Pesaro M, Rüdert A, Widmer F (2008). Resilience of the rhizosphere *Pseudomonas* and ammonia-oxidizing bacterial populations during phytoextraction of heavy metal polluted soil with poplar. *Environ. Microbiol.*, 10: 1433-1449.
- Holt J (1994). *Bergey's manual of systematic bacteriology*. 9th edn, Baltimore: Williams and Wilkins.
- Jin JR, Shen RX (1989). Study on microbial community in the soil of poplar stands and dominant species inhibiting *Dothiorella gregaria* (in Chinese). *J. Beijing Forestry Univ.*, 11: 79-84.
- Katamoto K (1986). Two new species of *Eudarlucella hyperparasitic* to *Botryosphaeria*. *Trans. Mycol. Soc. Jpn.*, 27: 11-16.
- Lee SJ, Park SY, Lee JJ, Yum DY, Koo BT, Lee JK (2002). Genes encoding the N-acyl homoserine lactone-degrading enzyme are widespread in many subspecies of *Bacillus thuringiensis*. *Appl. Environ. Microbiol.*, 68: 3919-3924.
- Moreno C, Romero J, Espejo RT (2002). Polymorphism in repeated 16S rRNA genes is a common property of type strains and environmental isolates of the genus *Vibrio*. *Microbiol.*, 148: 1233-1239.
- Morgan-Jones G, White JF (1987). Notes on Coelomycetes. II. Concerning the *Fusicoccum* anamorph of *Botryosphaeria ribis*. *Mycotaxon*, 30: 117-125.
- Romero D, de Vicente A, Rakotoaly RH, Dufour SE, Veening JW, Arrebola E, Cazorla FM, Kuiper OP (2007). The iturin and fengycin families of lipopeptides are key factors in antagonism of *Bacillus subtilis* towards *Podosphaera fusca*. *Mol. Plant Microbe Interact.*, 20: 430-440.
- Ryan RP, Germaine K, Franks A, Ryan DJ, Dowling DN (2008). Bacterial endophytes: recent developments and applications. *FEMS Microbiol. Lett.*, 278: 1-9.
- Stein T (2005). *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. *Mol. Microbiol.*, 56: 845-857.
- Stover AG, Wong PT (1999). Secretion localization, and antibacterial activity of *TasA*, A *Bacillus subtilis* spore-associated protein. *J. Bacteriol.*, 181: 1664-1672.
- Strange PN, Scott PR (2005). Plant disease: a threat to global food security. *Annu. Rev. Phytopathol.*, 43: 83-116.
- Ulrich K, Ulrich A, Ewald D (2008). Diversity of endophytic bacterial communities in poplar grown under field conditions. *FEMS Microbiol. Ecol.*, 63: 169-180.