

Full Length Research Paper

# Efficacy of seawater salinity on osmotic adjustment and solutes allocation in wheat (*Triticum aestivum*) flag leaf during grain filling

Heshmat S. Aldesuquy\*, Zakaria A. Baka, Omar A. El-Shehaby and Hanan E. Ghanem

Department of Botany, Faculty of Science, Mansoura University, Mansoura, Egypt.

Accepted 13 January, 2018

Two wheat (*Triticum aestivum* L.) cultivars (salt sensitive cultivar, Gemmieza-9 and salt resistant cultivar, Sids-1) subjected to different seawater salinity (10 and 25%). Osmotic pressure (OP), osmotic adjustment (OA) and solutes accumulation (TSS, TSN, proline, organic acids, glycerol and inorganic ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Cl}^-$ )) were quantified in flag leaf during grain-filling (14 and 21 days post-anthesis). Seawater salinity induced significant increase in osmotic pressure and the magnitude of increase was higher in Sids-1 than in Gemmieza-9. Furthermore, seawater concentrations caused noticeable increase in osmotic adjustment, organic solutes (TSS, TSN, proline, organic acids and glycerol) and inorganic ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Cl}^-$ ). On the other hand, clear reduction in  $\text{K}^+/\text{Na}^+$  ratio in the flag leaves of both cultivars was observed. The capacity of osmotic adjustment was greater in younger leaves than in older ones particularly with higher concentration (25%) in both cultivars. Moreover, the production of both organic and inorganic ions tended to be higher in Sids-1 than in Gemmieza-9. Gemmieza-9 appeared to be more sensitive than Sids-1. Osmotic pressure of flag leaf sap appeared to depend mainly on proline, TSN, TSS, organic acids, glycerol and ions content, where there is a positive correlation between osmotic pressure and all of them.

**Key words:** Wheat, seawater, osmotic adjustment, compatible solutes, glycerol.

## INTRODUCTION

Soil and water salinity have been considered a limiting factor to crop production in arid and semiarid regions of the world (Denden et al., 2005). Salinity effects on plants include two distinct types of stress: water stress, caused by the greater difficulty of water absorption, and ionic stress, related to the sodium ion effect on the diverse cellular functions, decreased nutrient absorption, enzyme activities, photosynthesis and metabolism (Zhu, 2001; Hajlaouia et al., 2006).

Plants have developed various combating mechanisms to survive with the deleterious effects of salt stress. Among these, osmotic adjustment (OA) is one of the strategies that have been a potential defense toward salt stress (Hajlaouia et al., 2010). This phenomenon is considered to be an important component of salinity tolerance mechanisms in plants (Neocleous and Vasilakakis, 2007) and also necessary to maintain water uptake from a saline soil (Ottow et al., 2005). Hence, OA allows water uptake, cell enlargement and plant growth during water stress associated with partial stomata opening allowing the  $\text{CO}_2$  assimilation at low water potentials that are otherwise inhibitory (Alves and Setter, 2004).

According to Blum et al. (1996), OA is usually defined as a decrease in cell sap osmotic potential resulting from a net increase in intracellular solutes rather than from a loss of cell water. The former may operate through the

\*Corresponding author. E-mail: HS-Aldesuquy@hotmail.com.

**Abbreviations:** OA, Osmotic adjustment; d, days post-anthesis; LSD, least significant difference; R, resistant; S, sensitive; SW, seawater; TSN, total soluble nitrogen; TSS, total soluble sugar

concentration accretion of inorganic and/or organic solutes (Ben Khaled et al., 2003). As a consequence, the cell's osmotic potential is diminished which in turn attracts water into the cell by tending to maintain turgor pressure (Pérez-Pérez et al., 2009). Furthermore, Munns (2005), compatible solutes like sugars, amino acids, organic acids and inorganic ions can contribute to this process as well as glycerol (Martinez et al., 2004, 2005) and fatty acids. Plants with increased concentrations of these compounds are expected to display increased salt tolerance (Taji et al., 2002).

The degree of OA also could be affected by the rate of stress intensity and most particularly by organ type and plant age (Alves and Setter, 2004). The occurrence of active OA can be established only if a net increase of solute concentration occurs (Silveira et al., 2009). Moreover, Ouda (2006) reported that many plants are able to partially compensate for low osmotic potential of the soil water by building up higher internal solute contents under saline conditions.

Sugars contributed up to 50% of the total osmotic potential in glycophytes subjected to saline conditions (Ashraf and Harris, 2004). Sugars accumulation in salinity stressed plants prevents structural and functional changes of membranes and destruction of soluble proteins. In accordance with these findings, Hichem et al. (2009) stated that, stressing two maize cultivars with various levels of salinity resulted in marked accumulation of sugars in roots and leaves of such stressed plants. Also, the accumulation of total soluble nitrogen (TSN) is another important adaptive response of plants to the lack of water and salt stress. Nitrogen-containing compounds are well documented to be involved as main osmolytes contributing in the plant osmotic regulation under salt stress. In this respect, Hassan et al. (2004) found that salinity stress increased TSN in fenugreek, mungbean and tomato plants.

Proline is one of the key osmolytes contributing toward osmotic adjustment (Soudry et al., 2005; Iqbal et al., 2008). Salinity increased markedly the proline content in different salt sensitive and tolerant species/cultivars with greater proline accumulation in salt tolerant ones, which is supposed to correlate with the adaptation to salinity (Ashraf and Harris, 2004; Mansour et al., 2005). In connection, salt stress remarkably enhanced proline accumulation in leaves of two rice cultivars differing in salinity tolerance and the rate of increase was more in the tolerant one, which implies the involvement of proline accumulation in osmotic adjustment during salinity (Demiral and Türkan, 2006). Srivastav et al. (2010) noticed that salt stress resulted in significant increase in proline content of mango leaves.

The accumulation of glycerol is another important adaptive response of plants and other organisms to the lack of water and saline stress. Glycerol has been shown to be a compatible solute for cells exposed to lowered water potential (Managbanag and Torzilli, 2002). In this connection, Torzilli (1997) found that, the salt stress

increases the cellular concentrations of glycerol in *Aureobasidium pullulans*.

Accumulation of inorganic solutes, such as cations ( $\text{Na}^+$  and  $\text{K}^+$ ) and the anion ( $\text{Cl}^-$ ), can also play a role independently or in combination with other mechanisms in maintaining the osmotic imbalance caused by the salt stress and influence the osmotic potential adjustment of plant cells (Bayuelo-Jimenez et al., 2003; Peng et al., 2004). Since the cytosolic and organellar machinery of glycophytes is equivalently  $\text{Na}^+$  and  $\text{Cl}^-$  sensitive, osmotic adjustment is achieved in these compartments by the accumulation of compatible osmolytes and osmo-protectants "which are small electrically neutral molecules that are nontoxic at molar concentrations and stabilize proteins and membranes against the denaturing effect of high concentrations of salts and other harmful solutes" (Munns, 2002).

The present study was undertaken to examine the effect of seawater salinity on osmotic adjustment and solutes allocation in wheat flag leaf during grain filling.

## MATERIALS AND METHODS

### Plant material and growth conditions

Pure strains of *Triticum aestivum* L. Gemmieza-9 (salt sensitive cultivar) and Sids-1 (salt resistant cultivar) were kindly supported by the Agricultural Research Center, Ministry of Agriculture, Giza, Egypt. For soaking experiment, a homogenous lot of *Triticum aestivum* L. (either sensitive or resistant cultivar) grains were selected. The grains were separately surface sterilized by soaking in 0.01 M  $\text{HgCl}_2$  solution for 3 min, then washed thoroughly with distilled water. The sterilized grains from each cultivar were drilled in plastic pots (25 cm in diameter) filled with 7 kg soil (clay: sand 2/1, v/v), where 15 grains were sown in each pot. The pots were then kept in a greenhouse at Botany Department, Faculty of Science, Mansoura University, Egypt. The plants were subjected to natural day/night conditions (min./max. air temperature and relative humidity were 15/25°C and 35/45%, respectively) at mid-day during the experimental period. The plants were irrigated to field capacity by tap water. After two weeks from sowing, thinning was started so that five uniform seedlings were left in each pot for the subsequent studies. The plants from each cultivar were divided into three sets. The 1<sup>st</sup> set was still irrigated with normal tap water serving as control, whereas the 2<sup>nd</sup> or 3<sup>rd</sup> ones were irrigated with 10 and 25% seawater receptively. Irrigation with seawater was applied after 30 days from sowing with a periodical soil washing (each two weeks) with tap water. After thinning and at heading, the plants received 36 kg N ha<sup>-1</sup> as urea and 25 kg P ha<sup>-1</sup> as superphosphate.

The chemical analyses of the employed seawater, collected from the Mediterranean Sea, revealed that it contains  $\text{Cl}^-$ , 21.6 Kg m<sup>-3</sup>;  $\text{Na}^+$ , 11.1 Kg m<sup>-3</sup>;  $\text{SO}_4^{2-}$ , 2.85 Kg m<sup>-3</sup>;  $\text{K}^+$ , 0.49 Kg m<sup>-3</sup> and  $\text{P}^{+3}$  16.6 µg dm<sup>-3</sup>. Its salinity was found to be 38.5 g kg<sup>-1</sup>; pH, 8.1 and EC, 47 mmhos cm<sup>-1</sup> (Aldesuquy and Baka, 1998). Samples were taken for measurements of osmotic pressure and other osmolytes during grain-filling ((14 and 21 days post-anthesis) (that is, 99 and 106 days after sowing)).

### Measurement of osmotic pressure (OP) and osmotic adjustment (OA)

The osmotic pressure of flag leaf sap was measured by the

cryoscopic method (Walter, 1949) and described by El-Sharkawi and Abdel-Rahman (1974). Osmotic adjustment (OA) was calculated as the difference in OP between salinized and control plants (Martinez-Ballesta et al., 2004)

#### **Determination of total soluble sugars**

Total soluble sugars was extracted and determined by anthrone method of Riazzi et al. (1985) as modified by Ibrahim (1999).

#### **Determination of the total soluble nitrogen**

The total soluble nitrogen was determined by the conventional semi micro-modification of Kjeldahl method (Pine, 1955).

#### **Estimation of proline**

The method adopted for estimation of proline was essentially that described by Snell and Snell (1954).

#### **Determination of keto acids**

Keto acids were determined according to the method adopted by Friedman and Haugen (1943).

#### **Determination of citric acid**

The method adapted for estimation of citric acid was essentially that described by Snell and Snell (1949).

#### **Estimation of glycerol**

Glycerol was estimated according to the method described and adopted by Mansour (1972).

#### **Determination of some mineral ions**

The extracts of the experimental plants were analyzed for the cations: Na<sup>+</sup>, K<sup>+</sup> and Ca<sup>+2</sup> Mg<sup>+2</sup> measured by flame emission spectrophotometry according to the method described by Chapman and Pratt (1978) and the anions Cl<sup>-</sup> chlorides were determined by the AgNO<sub>3</sub> titration method as described by Hansen and Munns (1988).

The sodium adsorption ratio (SAR) and potassium adsorption ratio (PAR) were calculated according to McKell and Goodin (1984) as:

$$SAR = \frac{Na^+}{\sqrt{\frac{Ca^{++} + Mg^{++}}{2}}}$$

$$PAR = \frac{K^+}{\sqrt{\frac{Ca^{++} + Mg^{++}}{2}}}$$

Where, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>++</sup> and Mg<sup>++</sup> refer to the concentrations of the designated cations.

#### **Statistical analysis**

A test for significant differences between means at P ≤ 0.05 was performed using least significant difference (LSD) test (Snedecor and Cochran, 1976). The correlation coefficients were estimated according to SPSS programme.

## **RESULTS**

### **Changes in osmotic pressure (OP) and osmotic adjustment (OA)**

Seawater stress induced a sharp increase (P ≤ 0.05) in OP and OA in flag leaf of both cultivars during grain filling as compared to control values (Figure 1). The resistant plants showed higher OP and OA values than the sensitive one under stress conditions. The difference, in terms of OP and OA, between cultivars was more conspicuous especially at higher seawater concentration (25%) where, the young leaves showed to have the maximum OA comparably to older ones particularly (25%). Generally, the Sids-1 cultivar exhibited more osmotic potential and osmotic adjustment.

### **Changes in total soluble sugar (TSS)**

The pattern of results in Figure 2 showed that, there is a noticeable increase in TSS in control and seawater-stressed plants from 14 to 21 days post-anthesis in both wheat cultivars. In relation to wheat cultivar, the resistant one had lower TSS values than the sensitive one. Seawater salinity induced a marked increase (P ≤ 0.05) in TSS in flag leaf of both cultivars during grain filling as compared to control values. It is clear that the resistant plants accumulated more TSS than the sensitive one under seawater-stress.

### **Changes in total soluble nitrogen (TSN)**

In relation to control values, seawater-stress induced a noticeable increase (P ≤ 0.05) in TSN from the younger leaves to the older leaves of both cultivars during anthesis stage, but the resistant plants accumulated more TSN than the sensitive one (Figure 3).


### **Changes in proline**

As compared to the control values, salinity stress caused apparent increase (P ≤ 0.05) in proline concentration in flag leaf of both cultivars during grain filling (Figure 4). Comparing both cultivars, under saline conditions, Sids-1 had more proline level than Gemmieza-9.


### **Changes in organic acids**

#### **Changes in keto-acids**


Perusal of the data shown in Figure 5 showed that, there is a marked decrease in keto-acids in control and seawater-stressed plants from 14 to 21 days post-anthesis in both wheat cultivars. In relation to control




**Figure 1.** Effect of different concentrations of seawater on OP and OA (atm) in flag leaf extract of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at  $P \leq 0.05$ .




**Figure 2.** Effect of different concentrations of seawater on TSS ( $\text{mg g}^{-1} \text{d wt}$ ) in flag leaf extract of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at  $P \leq 0.05$ .



**Figure 3.** Effect of different concentrations of seawater on TSN ( $\text{mg g}^{-1} \text{d wt}$ ) in flag leaf extract of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at  $P \leq 0.05$ .



**Figure 4.** Effect of different concentrations of seawater proline (mg g<sup>-1</sup> d wt) in flag leaf extract of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at P ≤ 0.05.



**Figure 5.** Effect of different concentrations of seawater on keto acids and citric acid (mg g<sup>-1</sup> d wt) in the extract of the flag leaf of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at P ≤ 0.05.

values, keto-acids accumulation in response to seawater-stress was more pronounced ( $P \leq 0.05$ ) in flag leaf of both cultivars during grain filling where the resistant plants accumulated more keto-acids than the sensitive ones (Figure 5).


#### Changes in citric acid

There is a tendency among control and seawater-stressed plants to a progressive increase in citric acid from 14 to 21 days post-anthesis in both wheat cultivars.

Seawater at all examined concentrations caused a marked increase ( $P \leq 0.05$ ) in citric acid in flag leaf of both cultivars at 14 and 21 days post-anthesis in comparing with control plants (Figure 5). Moreover, the resistant plants accumulated more citric acid than the sensitive one (Figure 5).

#### Changes in glycerol

The pattern of results in Figure 6 showed that, there is a noticeable increase in glycerol in control and seawater-



**Figure 6.** Effect of different concentrations of seawater on glycerol ( $\text{mg g}^{-1} \text{ d wt}$ ) in the extract of the flag leaf of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at  $P \leq 0.05$ .

stressed plants from 14 to 21 days post-anthesis in both wheat cultivars. In relation to control, seawater irrigation stimulated a progressive increase ( $P \leq 0.05$ ) in glycerol content in flag leaf of both cultivars during grain filling. The resistant cultivar accumulated more glycerol than the sensitive one under salt-stress. Almost, the increase was non-significantly at 10% in case of Gemmieza-9.

### Changes in ionic content

Perusal of the data shown in Figures 7 and 8 cleared that, there is a tendency among control and seawater-stressed plants to a progressive increase in ionic contents from 14 to 21 days post-anthesis in both wheat cultivars. In relation to control values, the two examined concentrations of seawater induced significant increase ( $P \leq 0.05$ ) in ions content ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{Cl}^-$ ) as well as  $\text{Na}^+$  adsorption ratio and  $\text{K}^+$  adsorption ratio and induced a noticeable decrease in  $\text{K}^+/\text{Na}^+$  ratio in flag leaf of both cultivars during anthesis stages (Figures 7 and 8).

In the majority of cases, the applied 10% dose of seawater revealed a non-significant increase on all the previous parameters (OP, TSS, TSN, keto-acids, citric acid,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$  adsorption ratio and  $\text{K}^+$  adsorption ratio) (in case of Gemmieza-9 in both stage except for  $\text{Na}^+$  and  $\text{Cl}^-$  ions, the increment in them was significantly in Gemmieza-9 and non-significantly in Sids-1. Hence, Gemmieza-9 depicted more susceptibility to salt toxicity in comparison to Sids-1 under saline conditions.


### DISCUSSION

Several reports are available that showed different strategies adopted to counteract the salinity effects (Ashraf, 2009; Türkan and Demiral, 2009) but the information on mineral nutrient status of plants and salinity tolerance is scarce (Khan et al., 2009b, 2010;

Khorshidi et al., 2009). Plants have developed various combating mechanisms to survive with the deleterious effects of salt stress. Among these, osmotic adjustment is one of the strategies that have been a potential defense toward seawater-salinity.

In the present study, seawater at the two used concentrations induced noticeable increase in osmotic pressure, some organic solutes (i.e. TSS, TSN, proline, organic acids and glycerol) and inorganic ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Cl}^-$ ) content as well as  $\text{Na}^+$  adsorption ratio and  $\text{K}^+$  adsorption ratio while induced a noticeable decrease in  $\text{K}^+/\text{Na}^+$  ratio in water extract of the flag leaf of both cultivars. Sids-1 accumulated more osmolytes than Gemmieza-9 (Figures 1-8). This increase in osmotic potential might be due to the increase in organic acids, TSS, TSN and inorganic osmolytes (i.e.  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Cl}^-$ ). Furthermore, osmotic pressure appeared to be positively correlated with organic osmolytes [TSS ( $r = 0.99, 0.94$ ); ( $r = 1.00, 0.93$ ), TSN ( $r = 0.99, 0.97$ ); ( $r = 0.99, 1.00$ ), proline ( $r = 0.99, 0.90$ ); ( $r = 0.99, 0.91$ ), keto-acids ( $r = 0.94, 0.99$ ); ( $r = 0.99, 0.99$ ), citric acid ( $r = 0.98, 0.95$ ); ( $r = 0.96, 0.92$ ) and glycerol ( $r = 0.99, 0.95$ ); ( $r = 0.99, 0.97$ )] as well as inorganic ions [( $\text{Na}^+$  ( $r = 1.00, 0.96$ ); ( $r = 0.98, 0.87$ ),  $\text{K}^+$  ( $r = 1.00, 0.91$ ); ( $r = 0.96, 0.92$ ),  $\text{Ca}^{2+}$  ( $r = 0.99, 0.93$ ); ( $r = 0.94, 0.95$ ),  $\text{Mg}^{2+}$  ( $r = 0.99, 0.93$ ); ( $r = 0.99, 0.96$ ),  $\text{Cl}^-$  ( $r = 0.98, 1.00$ ); ( $r = 0.98, 1.00$ )] for both Gemmieza-9 and Sids-1 during grain filling respectively (Table 1. In accordance to these results, Kusaka et al. (2005) reported that the observed increase in the osmotic potential of salt-stressed plants might be due to the accumulation of inorganic solutes (e.g.,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$  and  $\text{HPO}_4^-$ ), several organic components such as sucrose, glucose, quaternary ammonium compounds and amino acids.

Osmotic adjustment is a mechanism used for maintaining turgor and reducing the deleterious effects of salt stress on vegetative and reproductive tissue. Thus, salinity stress induced a marked increase in osmotic potential and osmotic adjustment of wheat flag leaf of both cultivars (Figure 1).




**Figure 7.** Effect of different concentrations of seawater on Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup> (mmol g<sup>-1</sup> d wt) in flag leaf extract of wheat cultivars during grain-filling (14 and 21 days post-anthesis). Vertical bars represent LSD at P ≤ 0.05.

These results were in agreement with those obtained by Ottow et al. (2005) who cleared that, by doubling NaCl concentration in the nutrient solution of *Populus euphratica* plants, these plants were able to adjust the osmotic pressure of leaves to levels just exceeding those of the nutrient solution, which is important to maintain water uptake and to prevent dehydration.

In comparing with control plants, soil drench with seawater at the two examined concentrations caused marked increase in TSS in flag leaf of both cultivars during grain filling with marked accumulation in Sids-1 comparing to Gemmieza-9 (Figure 2). In support, Hassanein et al. (2009) reported that various doses of NaCl, induced significant increases in soluble sugars in shoots of maize plants. Currently, sugar accumulation in

plants in response to seawater stress, is also quite well documented and confirming previous results reporting that soluble sugars are the main contributors to osmotic adjustment in stressed tissues (Xuana and Catherine, 2009; Gorai et al., 2010). In this respect, soluble carbohydrates (glucose, fructose, sucrose, fructans) accumulated under salt stress play a leading role in osmoprotection, osmotic adjustment, carbon storage and radical scavenging (Parida et al., 2002) and can be used as an indicator of salt tolerance (Juan et al., 2005; Almodares et al., 2008). Furthermore, Bartels and Sunkar (2005) suggested that in cotton leaves sugars act for osmotic adjustment and/or protect specific macromolecules and contribute to the stabilization of membrane structures, where sugars are thought to



**Figure 8.** Effect of different concentrations of seawater on Cl<sup>-</sup> (mmol g<sup>-1</sup> dwt), K<sup>+</sup>/Na<sup>+</sup> ratio, Na<sup>+</sup> adsorption ratio and K<sup>+</sup> adsorption ratio in flag leaf extract of wheat cultivars during grain-filling (14 & 21 days post-anthesis). Vertical bars represent LSD at P ≤ 0.05.

interact with polar head groups of phospholipids in membranes so that membrane fusion is prevented.

The nitrogen contents of plants growing under salt stress show great diversity (Mahmood et al., 2008).

**Table 1.** Correlation coefficient (r) between osmotic pressure, some osmolytes and ions of the two seawater-stressed wheat cultivars during grain-filling (14 & 21 days post-anthesis).

Sensitive			Resistant		
Variable	r		Variable	r	
	14 d	21 d		14 d	21 d
Na <sup>+</sup>	1.00	0.96	Na <sup>+</sup>	0.98	0.87
K <sup>+</sup>	1.00	0.91	K <sup>+</sup>	0.96	0.92
Ca <sup>2+</sup>	0.99	0.93	Ca <sup>2+</sup>	0.94	0.95
Mg <sup>2+</sup>	0.99	0.93	Mg <sup>2+</sup>	0.99	0.96
Cl <sup>-</sup>	0.98	1.00	Cl <sup>-</sup>	0.98	1.00
K <sup>+</sup> / Na <sup>+</sup> ratio	-0.99	-0.99	K <sup>+</sup> / Na <sup>+</sup> ratio	-0.92	-0.91
Na <sup>+</sup> adsorption ratio	1.00	0.95	Na <sup>+</sup> adsorption ratio	0.98	0.89
K <sup>+</sup> adsorption ratio	0.99	0.92	K <sup>+</sup> adsorption ratio	0.97	0.93
TSS	0.99	0.94	TSS	1.00	0.93
TSN	0.99	0.97	TSN	0.99	1.00
Proline	0.99	0.90	Proline	0.99	0.91
Keto-acids	0.94	0.99	Keto-acids	0.99	0.99
Citric acid	0.98	0.95	Citric acid	0.96	0.92
Glycerol	0.99	0.95	Glycerol	0.99	0.97

Wheat plants irrigated with seawater had higher TSN content in their flag leaf extract as compared to unstressed plants. Sids-1 was able to keep out its nitrogen content under seawater-stress at higher levels than Gemmieza-9, in order to increase their ability to withstand salinity stress (Figure 3). These results were in accordance with those of (Ehltling et al., 2007; Koyro et al., 2008) on Grey poplar and *Chenopodium quinoa* respectively.

The accumulation in TSN may result from a sharp increase in total free amino acids and total soluble proteins (Ibrahim, 2004). Likewise, this change in nitrogen content may be related to the inhibition of translocation from root to shoot, inhibition of protein synthesis or the increase in protease activity (Khalil and Mandurah, 1990). Moreover, this increase in the soluble nitrogen compounds is of important in plant osmoregulation in response to salinity stress.

The pattern of proline accumulation in the flag leaves of the two wheat cultivars showed gradual increase with increasing salinity level however Sids-1 had higher proline accumulation than Gemmieza-9 (Figure 4). Sids-1 is better performing than Gemmieza-9 under salinity stress. Similar observations were recorded by many workers (Pagter et al., 2009; Khan et al., 2009a; Salah et al., 2011) in different crops. Proline accumulation is one of the common characteristics in many plants exposed to salt stress (Ashraf and Harris, 2004). Therefore, proline could be used as is a good parameter for the evaluation of tolerance or sensitivity of plants to stress (Misra and Gupta, 2004; Shi and Wang, 2005). Therefore, proline accumulation may contribute to osmotic adjustment at the cellular level (Mahajan and Tuteja, 2005; Demiral and

Türkan, 2006). In this respect, proline accumulation could be a protective response not only due to the its osmoprotectant role that prevents salinity-induced water deficit but also for its hydroxyl scavenger, stabilization of membrane and protein structure, a sink of carbon and nitrogen for stress recovery, and buffering cellular redox potential under stress (Matysik et al., 2002; Khedr et al., 2003; Lee et al., 2008). Also, as an alternative to direct ROS scavenging feature, proline can protect and stabilize ROS scavenging enzymes and activate alternative detoxification pathways (Szabados and Savouré, 2009).

Organic acids content (mainly citric and keto-acids) were increased in flag leaf sap of seawater-stressed plants during grain filling with higher accumulation in Sids-1 (Figure 5). These results were in a good conformity with those obtained by many authors Shi and Sheng (2005) and Guo et al. (2009). In this respect, organic acids not only an important organic osmotic regulator but also an important negative charge contributor, playing important roles in ionic balance and pH adjustment. Additionally, they might reflect an adaptation trait of wheat plant in response to seawater irrigation. It is well known that organic acids play a major role in the salt stress adaptation of plants (Gilbert et al., 1998). The increase in citric acid content might be explained on the fact that, the role of citric acid decarboxylation to increase CO<sub>2</sub> concentration in mesophyll during the day, while night-time citric acid accumulation increases the buffer capacity of the vacuoles (Franco et al., 1992).

In comparing with control plants, seawater irrigation caused a marked increase in glycerol content of flag leaf of both cultivars during grain filling with marked accumulation in Sids-1 than Gemmieza-9 (Figure 6).

These results were also in line with those obtained by Ben-Amotz and Avron (1973) and Frank and Wegmann (1974), who reported that cells tended to accumulate high concentrations of glycerol with increasing the external salinity level in order to increasing the internal cell osmotic pressure. Such increment would improve the performance of the cell at high salinity by preventing leakage of the compatible solute out of the cell and diffusion of potentially harmful ions into the cell Elenkov et al. (1996). Glycerol has been shown to be a compatible solute involves in osmotic adjustment in cells exposed to lowered water potential (Martinez et al., 2004, 2005). Moreover, Torzilli (1997) suggests that increases in the cellular concentrations of glycerol contribute to the acquisition of stress tolerance. Also, glycerol was required to balance the osmotic potential of  $\text{Na}^+$  and  $\text{Cl}^-$  (Flowers and Colmer, 2008). The effect of hypertonic concentration of NaCl has been attributed to the formation of glycerol as osmotic agent to facilitate the retention of cellular water (Berry and Brown, 1987) and maintain enzyme activity (Borowitzka and Brown, 1974).

Excess or deficiency of any mineral nutrient is crucial for the reason that the plant growth depends on supply of inorganic nutrients (Marschner, 1995). Although organic compounds are the major compounds of osmoregulation in plant cells during water deficit stress, inorganic ions would also contribute to the osmotic adjustment (Guo et al., 2009). In addition, synthesis and accumulation of organic solutes consume more energy than uptake of inorganic ions (Xu and Yu, 1990). Generally, the obtained results in Figures 7 and 8 showed that seawater at all examined concentrations induced marked increase in  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Cl}^-$  contents as well as  $\text{Na}^+$  adsorption ratio and  $\text{K}^+$  adsorption ratio. A clear reduction in  $\text{K}^+/\text{Na}^+$  ratio was observed in wheat flag leaf of both cultivars during grain filling. These results were in agreement with those obtained by Jin et al. (2007) who suggested that seawater salinity increased the level of inorganic cations in both root and shoot of *Aloe vera* plants.  $\text{Na}^+$  and  $\text{Cl}^-$  contents increased due to salinity in the two wheat genotypes however genotypes Gemmieza-9 maintained the higher leaf  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations than Sids-1 (Figures 7 and 8). These results were in agreement with those obtained by Patakas et al. (2002), who observed an increase in the  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations in grapevine leaves under water stress. Moreover, Silva et al. (2010) cleared that a significant increase in both  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations in *Jatropha curcas* plants treated with 50 mM NaCl.

$\text{Na}^+$  in  $\text{C}_4$  species such as wheat is related to its involvement in photosynthesis mainly in the maintenance of mesophyll chloroplast structure, mainly in relation to granal stacking (Brownell and Bieligi, 1996). In addition, under saline conditions  $\text{Na}^+$  influx across the plasma lemma to the vacuole might play a major role in permitting turgor maintenance, and this accumulation of  $\text{Na}^+$  inside the vacuoles reduce its toxic levels in cytosol

and increase the vacuolar osmotic potential with the concomitant generation of a more negative water potential that favors water uptake by the cell and better tissue water retention under high salinity levels (Tawfik et al., 2006). Additionally, the greatest accumulation of  $\text{Na}^+$  by plants at high salt concentration may be attributed to the damage of the protoplasm of plant cells and as a result, the selective salt absorption is replaced by passive absorption which causes abnormal accumulation of salts in plant organs (Kader and Lindberg, 2005). To date, very little is known about the toxic effects of  $\text{Cl}^-$  ions in the plant cell (Natasha and Stephen, 2010). However, in the leaves of citrus plants,  $\text{Cl}^-$  ions are more toxic than  $\text{Na}^+$  ions but responsible mechanisms are yet unknown (Storey and Walker, 1999). It is also reported that sensitivity of some crops to salinity is due to the inability to keep  $\text{Na}^+$  and  $\text{Cl}^-$  out of transpiration streams (Gorham et al., 1990). Since  $\text{Na}^+$  and  $\text{Cl}^-$  are toxic elements whose higher concentration disturbs the different metabolic activities, Sids-1 which successes in retaining  $\text{Na}^+$  and  $\text{Cl}^-$  in the root was more tolerant than Gemmieza-9 which translocated maximum  $\text{Na}^+$  and  $\text{Cl}^-$  in leaves.

Data in Figure 7 showed that in response to salinity stress,  $\text{K}^+$  concentration increased in flag leaves of both cultivars with a higher rate in Sids-1. This is consistent with findings reported on *Medicago truncatula* and *Medicago laciniata* populations (Yousfi et al., 2010).  $\text{K}^+$  is known to function in osmotic adjustment in the guard cell controlling the stomatal movements and thus  $\text{CO}_2$  assimilation in photosynthesis (Degl'Innocenti et al., 2009; Cha-um et al., 2010). Similarly, changes in the potassium content may contribute substantially to osmoregulation (Shabala and Cuin, 2007) and may occur in concert with changes in sugars and amino acids (Pérez-Pérez et al., 2009).

Seawater stress caused a decrease in  $\text{K}^+/\text{Na}^+$  ratio in wheat flag leaves of both wheat cultivars with a higher magnitude of response in Gemmieza-9 (Figure 8). This reduction in  $\text{K}^+/\text{Na}^+$  ratio in wheat leaves with increasing salinity level might be due to competition of  $\text{Na}^+$  with  $\text{K}^+$  where this competition could be at uptake level and/or transport level. Since maintenance of a high cytosolic  $\text{K}^+/\text{Na}^+$  ratio is a key feature of plant salt tolerance, as indicated by Cuin et al. (2008) and Kronzucker et al. (2008). Additionally, intracellular  $\text{K}^+$  and  $\text{Na}^+$  homeostasis bears importance for the activities of many cytosolic enzymes, maintaining membrane potential and a suitable osmoticum for cell volume regulation (Shabala and Cuin, 2007; Munns and Tester, 2008). These results are supported by Grewal (2010) who observed that increasing levels of subsoil NaCl salinity had significant depressing effect on leaves  $\text{K}^+/\text{Na}^+$  ratio of wheat, barley, canola and chickpea.

Perusal data showed that  $\text{Ca}^{2+}$  content was increased to some extent in flag leaf extract of both wheat cultivars during grain filling with more increment in Sids-1 (Figure 7). These results were in a harmony with those obtained

by Hessini et al. (2009) who observed that,  $\text{Ca}^{2+}$  presented a significant augment at the mild stress level in *Spartina alterniflora*. The increase in  $\text{Ca}^{2+}$  level as a result of seawater irrigation may enhance the tolerance of wheat plants to salinity stress since  $\text{Ca}^{2+}$  is a non-toxic inorganic nutrient and has a function of detoxification under saline medium (Izzo et al., 2008). In this regard,  $\text{Ca}^{2+}$  enhanced salt tolerance in barley (Huang and Redman, 1995) and sorghum (Bernstein et al., 1993).

Moreover, plants respond directly and specifically to  $\text{Na}^+$  within seconds by increasing cytosolic  $\text{Ca}^{2+}$  (Munns and Tester, 2008). In fact,  $\text{Ca}^{2+}$  is known to play an important role in processes that preserve the structural and functional integrity of plant membranes, regulate ion transport and control activities of cell wall enzymes (Rengel, 1992). Moreover, many researches showed that calcium signals decoding elements are involved in ABA-induced stomatal closure and plant adaptation to salt and other abiotic stresses and some new studies show that  $\text{Ca}^{2+}$  is dissolved in water in the apoplast and transported primarily from root to shoot through the transpiration stream (Song et al., 2008).

In conclusion, genotype Sids-1 appears to be more tolerant to seawater salinity than Gemmieza-9 since it able to maintain less amounts of  $\text{Na}^+$  and  $\text{Cl}^-$  and higher amounts of organic solutes (TSS, TSN, proline, organic acids, glycerol and inorganic ions ( $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{K}^+/\text{Na}^+$  ratio) in their flag leaves. In addition, this study provides scarce evidence supporting the hypothesis that OA plays a preponderant role in the resistance to seawater stress in both wheat cultivars. Also, the degree of OA also affected by the rate of stress intensity. Results of this research suggest also that OA could be a part of the salt tolerance mechanisms developed by wheat and could be exploited in breeding programs for improved salt stress tolerance.

## REFERENCES

- Aldesuquy HS, Baka ZAM (1998). Interactive effects of seawater and plant hormones on the pigments content and chloroplast ultrastructure of wheat flag leaf. Proceedings, Sixth Egyptian Botanical Conference, Cairo University, Giza. November 24- 26, 1998. 1: 51- 46.
- Almodares A, Ham MR, Dosti B (2008). The effects of salt stress on growth parameters and carbohydrates contents in sweet sorghum. Res. J. Sci., 2: 298- 304.
- Alves AAC, Setter TL (2004). Abscisic acid accumulation and osmotic adjustment in cassava under water deficit. Environ. and Exp. Bot., 51: 259- 271
- Ashraf M (2009). Biotechnological approach of improving plant salt tolerance using antioxidants as markers. J. Advan. Biotechnol., 27: 84- 93.
- Ashraf M, Harris PJC (2004). Potential biochemical indicators of salinity tolerance in plant. Plant Sci., 166: 3- 16
- Bartels D, Sunkar R (2005). Drought and salt tolerance in plants. CRC Critical Rev. Plant Sci., 24: 23- 58.
- Bayuelo-Jimenez JS, Debouck DG, Lynch JP (2003). Growth, gas exchange, water relations, and ion composition of *Phaseolus* species grown under saline conditions. Res. J. Field Crop 80: 207- 222.
- Ben-Amotz A, Avron M (1973). The role of glycerol in the osmotic regulation of the halophilic alga (*Dunaliella parva*). J. Plant Physiol., 51: 875- 878.
- Ben Khaled L., Morte-Gomez A, Honrubia M, Oihabi A (2003). Effet du stress salin en milieu hydroponique sur le tréfle inoculé par le Rhizobium. Agronomie 23: 553- 560.
- Bernstein N, Lauchli A, Silk W (1993). Kinematics and dynamics of sorghum (*Sorghum bicolor* L.) leaf development at various NaCl salinities. Plant Physiol., 103: 1107- 1114.
- Berry DR, Brown C (1987). Glycerol formation in yeast biotechnology. (eds. Berry, D.R.; Russell, I. and Stewart, G.G.), Allen Nd Unwin, London, 185- 186.
- Blum A, Munns R, Passioura JB, Turner NC, Sharp RE, Boyer JS, Nguyen HT, Hsiao TC (1996). Letters to the editor genetically engineered plants resistant to soil drying and salt stress: how to interpret osmotic relations? Plant Physiol., 110: 1051- 1053.
- Borowitzka LJ, Brown AD (1974). The salt relations of marine and halophilic species of unicellular green algae, Dunaliella: The role of glycerol as a compatible solute. Arch. Microbiol., 96: 37- 52.
- Brownell PF, Bieligi LM (1996). The role of sodium in the conversion of pyruvate to phosphoenolpyruvate in the mesophyll chloroplasts of C3 plants. Aust. J. Plant Physiol., 23: 171- 177.
- Chapman HD, Pratt PF (1978). Methods of analysis for soils, plants and waters. University of California, Division of Agriculture and Science. pp 207-230
- Cha-um S, Siringamb K, Juntawongb N, Kirdmanee C (2010). Water relations, pigment stabilization, photosynthetic abilities and growth improvement in salt stressed rice plants treated with exogenous potassium nitrate application. Inter. J. Plant Prod., 4: 187- 198.
- Cuin TA., Betts SA, Chalmandrier R, Shabala S (2008). A root's ability to retain  $\text{K}^+$  correlates with salt tolerance in wheat. J. Exp. Bot., 59: 2697- 2706.
- Degl'Innocenti E, Hafsi C, Guidi L, Navari-Izzo F (2009). The effect of salinity on photosynthetic activity in potassium-deficient barley species. J. Plant Physiol., 166: 1968- 1981.
- Demiral T, Turkan I (2006). Exogenous glycinebetaine affects growth and proline accumulation and retards senescence in two rice cultivars under NaCl stress. Environ. Exp. Bot., 56: 72- 79.
- Denden M, Bettaieb T, Salhi A, Mathlouthi M (2005). Effet de la salinite sur la fluorescence chlorophyllienne, la teneur en proline et la production florale de trois especes ornementales. Tropicultura 23: 220-225
- Ehling B., Dluzniewska P, Dietrich H, Selle A (2007). Interaction of nitrogen nutrition and salinity in Grey poplar (*Populus tremula x alba*). Plant Cell Environ., 12: 1- 16.
- Elenkov L, Stefanov K, Dimitrova-Konaklieva S, Popov S (1996). Effect of salinity on lipid composition of *Cladophora vagabunda*. Phytochem., 42: 39- 44.
- El-Sharkawi HM, Abdel Rahman AA (1974). Response of olive and almond orchards to partial irrigation under dry farming practices in semi arid regions. Plant soil water relations in olive during the growing season. Plant Soil 41: 13- 32.
- Flowers TJ, Colmer TD (2008). Salinity tolerance in halophytes. New Phytolo., 179: 945- 963.
- Franco AC, Ball E, Lutge U (1992). Differential effects of drought and light levels on accumulation of citric acid and malic acid during CAM in *Clusia*. Plant Cell and Environ., 15: 821- 829.
- Frank G, Wegmann K (1974). Physiology and biochemistry of glycerol biosynthesis in *Dunaliella*. Biologisches Zentralblatt 93: 707- 723.
- Friedman TE, Haugen GE (1943). Pyruvic acid. II. The determination of keto acids in blood and urine. J. Biol. Chem., 147: 415- 442.
- Gilbert GA, Gadush MV, Wilso C, Madore MA (1998). Amino acid accumulation in sink and source tissues of *Coleus blumei* during salinity stress. J. Exp. Bot., 49: 107- 114.
- Gorai M, Ennajeh M, Khemira H, Neffati M (2010). Combined effect of NaCl-salinity and hypoxia on growth, photosynthesis, water relations and solute accumulation in *Phragmites australis* plants Mustapha. Flora 205: 462- 470.
- Gorham J, Wyn Jones RG, Bristol A (1990). Partial characterization of the trait for enhanced  $\text{K}^+/\text{Na}^+$  discrimination in the D-genome of wheat. Planta, 180: 590- 597.
- Grewal HS (2010). Water uptake, water use efficiency, plant growth and ionic balance of wheat, barley, canola and chickpea plants on a Sodic vertosol with variable subsoil NaCl salinity. Agric Water

- Manage., 97: 148- 156.
- Guo LQ, Shi DC, Wang DL (2009). The key physiological response to alkali stress by the alkali-resistant halophyte *Puccinellia tenuiflora* is the accumulation of large quantities of organic acids and into the rhizosphere. *J. Agronom. Crop Sci.*, 196: 123- 135.
- Hajlaouia H, Denden M, Bousslama M (2006). Effet du chlorure de sodium sur les critères morpho-physiologiques et productifs du pois chiche (*Cicer arietinum* L.). *Ann. INRGREF* 8: 171- 187.
- Hajlaouia H, El Ayebe N, Garrecc JP, Dendend M (2010). Differential effects of salt stress on osmotic adjustment and solutes allocation on the basis of root and leaf tissue senescence of two silage maize (*Zea mays* L.) varieties. *Industrial Crops Prod.*, 31: 122- 130.
- Hansen EM, Munns DN (1988): Effect of CaSO<sub>4</sub> and NaCl on mineral content of *Leucaena leucocephala*. *Plant Soil* 107: 101- 105.
- Hassan NM, Serag MS, El-Feky FM (2004). Changes in nitrogen content and protein profiles following in vitro selection of NaCl resistant mungbean and tomato. *Acta Physiol. Plant* 26: 165- 175.
- Hassanein RA, Bassuony FM, Baraka DM, Khalil RR (2009). Physiological effects of nicotinamide and ascorbic acid on *Zea mays* plant grown under salinity stress. I. Changes in growth, some relevant metabolic activities and oxidative defense systems. *Res. J. Agric. Biol. Sci.*, 5: 72- 81.
- Hessini K, Martínez JP, Gandoura M, Albouchib A, Soltania A, Abdelya C (2009). Effect of water stress on growth, osmotic adjustment, cell wall elasticity and water-use efficiency in *Spartina alterniflora*. *Environ. and Exp. Bot.*, 67: 312- 319.
- Hichem H, Mounir D, Naceur EA (2009). Differential responses of two maize (*Zea mays* L.) varieties to salt stress: Changes in polyphenols composition of foliage and oxidative damages. *Indust. Crops Prod.*, 30: 144- 151.
- Huang J, Redman RE (1995). Responses of growth, morphology and anatomy to salinity and calcium supply in cultivated and wild barley. *Canad. J. Bot.*, 73: 1859- 1866.
- Ibrahim AH (2004). Efficacy of exogenous glycine betaine application on sorghum plants grown under salinity stress. *Acta Bot. Hung.*, 43: 307- 318.
- Ibrahim IF (1999). Physiological effects of some uriede compounds on cultured *Phaseolus vulgaris* seedlings in vivo and in vitro. M. Sc. Thesis, Faculty of Science, Mansoura University, Egypt.
- Iqbal N, Ashraf M, Ashraf MY (2008). Glycinebetaine, an osmolyte of interest to improve water stress tolerance in sunflower (*Helianthus annuus* L.): water relations and yield. *South Afri. J. Bot.*, 74: 274-281.
- Izzo R, Incerti A, Bertolla C (2008). Seawater irrigation: Effects on growth and nutrient uptake of sunflower plants. In: *Biosaline Agric. High Salinity Tolerance*, 61- 69.
- Jin ZM, Wang CH, Liu ZP, Gong WJ (2007). Physiological and ecological characters studies on *Aloe vera* under soil salinity and seawater irrigation. *Processes Biochem.*, 42: 710- 714.
- Juan M, Rivero RM, Romero L, Ruiz JM (2005). Evaluation of some nutritional and biochemical indicators in selecting salt-resistant tomato cultivars. *Environ. Exp. Bot.*, 54: 193- 201.
- Kader MA, Lindberg S (2005). Uptake of sodium in protoplasts of salt-sensitive and salt-tolerant cultivars of rice (*Oryza sativa* L.) determined by the fluorescent dye SBFI. *J. Exp. Bot.*, 56: 3149- 3158.
- Khalil S, Mandurah HM (1990). Effects of water stress deficiency and kinetin on growth and nitrogen metabolism of cowpea plants. *J. Agron. Crop Sci.*, 164: 93- 99.
- Khan A, Nazar R, Anjum NA (2009b). Growth, photosynthesis and antioxidant metabolism in mustard (*Brassica juncea* L.) cultivars differing in ATP-sulfurylase activity under salinity stress. *Sci. Hortic.*, 122: 455- 460.
- Khan MA, Shirazi MU, Mujtaba SM, Slam E, Mumtaz S, Shereen A, Ansari RU, Ashraf YM (2009a). Role of proline, K<sup>+</sup>/Na<sup>+</sup> ratio and chlorophyll content in salt tolerance of wheat (*Triticum aestivum* L.). *Pakist. J. Bot.*, 41: 633- 638.
- Khan NA, Syeed S, Masood A, Nazar R, Iqbal N (2010). Application of salicylic acid increases contents of nutrients and antioxidative metabolism in mungbean and alleviates adverse effects of salinity stress. *Inter. J. Plant Biol.*, 1: 1-8.
- Khedr AHA, Abbas MA, Wahid AAA, Quick WP, Abogadallah GM (2003). Proline induces the expression of salt-stress responsive proteins and may improve the adaptation of (*Pancratium maritimum* L.) to salt-stress. *J. Exp. Bot.*, 54: 2553- 2562.
- Khorshidi MB, Yarnia M, Hassanpanah D (2009). Salinity effect on nutrients accumulation in alfalfa shoots in hydroponic condition. *J. Food Agric. Environ.*, 7: 787- 790.
- Koyro HW, Lieth H, Said S. (2008). Salt tolerance of *Chenopodium quinoa* Willd, grains of the Andes: Influence of salinity on biomass production, yield, composition of reserves in the seeds. water and solute. In: *Mangroves and Halophytes: Restoration and Utilisation. Part III*, 133- 145.
- Kronzucker H, Szczerba MW, Schulze LM, Britto DT (2008). Non-reciprocal interactions between K<sup>+</sup> and Na<sup>+</sup> ions in barley (*Hordeum vulgare* L.). *J Exp. Bot.*, 59: 2973- 2801.
- Kusaka, M, Lalusin AG, Fujimura T (2005). The maintenance of growth and turgor in pearl millet (*Pennisetum glaucum* L.) cultivars with different root structures and osmo-regulation under drought stress. *Plant Sci.*, 168: 1- 14.
- Lee VG, Carrow RN, Duncan RR, Eiteman MA, Mark WR (2008). Synthesis of organic osmolytes and salt tolerance mechanisms in *Paspalum*. *Environ. Exp. Bot.*, 63: 19- 27.
- Mahajan S, Tuteja N (2005). Cold, salinity and drought stresses. An overview: *Arch. Biochem. Bioph.*, 444: 139-158.
- Mahmood A, Athar M, Qadir R, Mahmood N (2008). Effect of NaCl salinity on growth, nodulation and total nitrogen content in *Sesbania sesban* (L.). *Agric. Conspetctus Scientificus*, 73: 137- 141.
- Managbanag JR, Torzilli AP (2002). An analysis of trehalose, glycerol, and mannitol accumulation during heat and salt stress in a salt marsh isolate of *Aureobasidium pullulans*. *Mycology*, 94, 384- 391.
- Mansour FA (1972). Physiology and biological studies on some local actinomycetes with special references to streptomycin production. M. Sc. Thesis. Cairo University, Cairo, Egypt.
- Mansour MMF, Salama KHA, Ali FZM, Abou Hadid AF (2005). Cell and plant responses to NaCl in (*Zea mays* L.) cultivars differing in salt tolerance. *General and Applied Plant Physiology*, 31: 29- 41.
- Marschner H (1995). *Mineral Nutrition of Higher Plants*, 2nd ed. Academic Press, San Diego, New York.
- Martínez JP, Kinet JM, Bajji M, Lutts S (2005). NaCl alleviates polyethylene glycol induced water stress in the halophyte species (*Atriplex halimus* L.). *J. Exp. Bot.*, 56: 2421- 2431.
- Martínez JP, Lutts S, Schanck A, Bajji M, Kinet, JM (2004). Is osmotic adjustment required for water stress resistance in the mediterranean shrub (*Atriplex halimus* L.)? *J Plant Physiol.*, 161: 1041- 1051.
- Martínez-Ballesta MC, Martínez V, Carvajal M (2004). Osmotic adjustment, water relations and gas exchange in pepper plants grown under NaCl or KCl. *Environ Exp. Bot.*, 52; 161-174
- McKell CM, Goodin JK (1984). A brief overview of the saline lands of the United States. Research and development seminar on forage and fuel production from salt-affected Wasteland, Western Australia.
- Matysik J, Alia B, Bhalu B, Mohanty P (2002). Molecular mechanisms of quenching of reactive oxygen species by proline under stress in plants. *Curr. Sci.*, 82: 525- 532.
- Misra AN, Gupta AK (2004). Effect of salt stress on proline metabolism in two high- yielding genotypes of green gram. *Plant Sci.*, 169: 331- 339.
- Munns R (2005). Genes and salt tolerance: Bringing them together. *New Phytologist.*, 167: 645- 663.
- Munns R, Tester M (2008). Mechanisms of salinity tolerance. *Ann. Rev. Plant Biol.*, 59: 651- 681.
- Natasha LT, Stephen DT (2010). Mechanisms of Cl<sup>-</sup> transport contributing to salt tolerance. *Plant Cell and Environ.*, 33: 566- 589.
- Neocleous D, Vasilakakis M (2007). Effects of NaCl stress on red raspberry (*Rubus idaeus* L. and *Autumn Bliss* L.). *Scientia Horticult.*, 112: 282- 289.
- Ottow EA, Brinker M, Teichmann T, Fritz E, Kaiser W, Brosché M, Kangasjärvi J, Jiang X and Polle A (2005). *Populus euphratica* displays apoplastic sodium accumulation, osmotic adjustment by decreases in calcium and soluble carbohydrates, and develops leaf succulence under salt stress. *Plant Physiol.*, 139: 1762- 1772.
- Ouda HAS (2006). Predicting the effect of water and salinity stresses on wheat yield and water needs. *J. Appl. Sci. Res.*, 2: 746- 750.
- Pagter M, Bragato C, Malagoli M, Brix H (2009). Osmotic and ionic effects of NaCl and Na<sub>2</sub>SO<sub>4</sub> salinity on *Phragmites australis*. *Aquatic*

Bot., 90: 43- 51.

- Parida AK, Das AB, Das P (2002). NaCl stress causes changes in photosynthetic pigments, proteins and other metabolic components in the leaves of a true mangrove (*Bruguiera parviflora*) in hydroponic cultures. *J. Plant Biol.*, 45: 28- 36.
- Patakas A, Nikolaou N, Zioziou E, Radoglou K, Noitsakis B (2002). The role of organic solute and ion accumulation in osmotic adjustment in drought-stressed grapevines. *Plant Sci.*, 163: 361- 367.
- Peng YH, Zhu YF, Mao YQ, Wang SW, Su WA, Tang ZC (2004). Alkali grass resists salt stress through high [K<sup>+</sup>] and an endodermis barrier to Na<sup>+</sup>. *J. Exp. Bot.*, 55: 939- 949.
- Pérez-Pérez JG, Robles JM, Tovar JC, Botía P (2009). Response to drought and salt stress of lemon 'Fino 49' under field conditions: water relations, osmotic adjustment and gas exchange. *Scientia Horticult.*, 122: 83- 90.
- Pine NW (1955). Proteins. In: Modern methods of plant analysis. (Peack, K. and Tracey, M. V., eds) IV, 23, Springer-Verlag, Berlin.
- Rengel Z (1992). The role of calcium in salt toxicity. *Plant Cell Environ.*, 15: 625- 632.
- Riazi A, Matsuda K, Arslan A (1985). Water stress induced changes in concentrations of proline and other solutes in growing regions of young barely leaves. *J. Exp. Bot.*, 36: 1716-1725.
- Salah BI, Slatni T, Gruber M, Messedia D, Gandoura M, Benzartia M, Haoualac R, Zribi K, Ben Hamed K, Perez-Alfoceae F, Abdelya C (2011). Relationship between symbiotic nitrogen fixation, sucrose synthesis and anti-oxidant activities in source leaves of two *Medicago ciliaris* lines cultivated under salt stress. *Environ. Exp. Bot.*, 70: 166- 173.
- Shabala S, Cui TA (2007). Potassium transport and plant salt tolerance. *Physiol. Plant* 133: 651- 669.
- Shi D, Sheng Y (2005). Effect of various salt-alkaline mixed stress conditions on sunflower seedlings and analysis of their stress factors. *Environ. Exp. Bot.*, 54: 8- 21.
- Shi DC, Wang DL (2005). Effects of various salt-alkali mixed stresses on *Aneurolepidium chinense* (Trin.). *Plant Soil*, 271: 15- 26.
- Silva EN, Ribeiro RV, Ferreira-Silva SL, Viégas RA, Silveira JAG (2010). Comparative effects of salinity and water stress on photosynthesis, water relations and growth of *Jatropha curcas* plants. *J. Arid Environ.*, 74: 1130- 1137.
- Silveira JA, Lima JPMS, Viégas RA (2009). Roots and leaves display contrasting osmotic adjustment mechanisms in response to NaCl-salinity in *Atriplex numularia*. *Environ. Exp. Bot.*, 66: 1- 8.
- Snedecor GW, Cochran WG (1976). *Statistical Methods*. 6th ed. Oxford IBH Publishing Co. New Delhi. pp. 217-235
- Snell FD, Snell CT (1949). *Colorimetric methods of analysis*. Volume II. D. Van Nostrand Co. Inc., New York. Pp. xxiii + 766
- Snell FD, Snell CT (1954). *Colorimetric methods of analysis*. Volume IV. D. Van Nostrand Co. Inc., New York. pp xxiii + 786
- Srivastava M, Kishora A, Dahujab A, Sharmac ARR (2010). Effect of paclobutrazol and salinity on ion leakage, proline content and activities of antioxidant enzymes in mango (*Mangifera indica* L.). *Scientia Horticult.*, 125: 785- 788.
- Song WY, Zheng-Bin Z, Shao HB, Guo Z L, Cao HX, Zhao HB, Fu ZY, Hu XJ (2008). Relationship between calcium decoding elements and plant abiotic-stress resistance. *Inter. J. Biol. Sci.*, 4: 116- 125.
- Soudry E, Ulitzur S, Gepstein S (2005). Accumulation and remobilization of amino acids during senescence of detached and attached leaves in planta analysis of tryptophan levels by recombinant luminiscent bacteria. *J. Exp. Bot.*, 56:695-702.
- Storey R, Walker RR (1999). Citrus and salinity. *Scientia Horticult.*, 78: 39- 81.
- Szabados L, Savouré A (2009). Proline: a multifunctional amino acid. *Trends Plant Sci.*, 15: 89- 97.
- Taji T, Ohsumi C, Iuchi S, Seki M, Kasuga M, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K (2002). Important roles of drought- and cold-inducible genes for galactinol synthase in stress tolerance in *Arabidopsis thaliana*. *Plant J.*, 29: 417- 426.
- Tawfik MM, Bahr AA, Salem AKM (2006). Response of kallar grass (*Leptochloa fusca* L.) to biofertilizer inoculation under different levels of seawater irrigation. *Res. J. Appl. Sci.*, 2: 1203- 1211.
- Torzilli AP (1997). Tolerance to high temperature and salt stress by a salt marsh isolate of *Aureobasidium pullulans*. *Mycology*, 98: 786- 792.
- Türkan I, Demiral T (2009). Recent developments in understanding salinity tolerance. *Environ and Exp. Bot.*, 67: 2- 6.
- Walter H (1949). *Gundlagen der flenzen verliertung*. Eintubring, in di pflanzengeographie- fur studierends der hocholen, standorstlehre: Stuttgart Ulmer. pp. 236.
- Xu YL, Yu SW (1990). Energy consumption in plant adapting to salinity adverse habitat. *Acta Phytophysiol.*, 6: 70- 73.
- Xuana L, Catherine G (2009) Accumulation of chiro-inositol and other non-structural carbohydrates in *Limonium* species in response to saline irrigation waters. *J. Amer. Soc. Horticult. Sci.*, 134: 329- 336.
- Yousfi N, Slama I, Ghnaya T, Savouré A, Abdely C (2010). Effects of water deficit stress on growth, water relations and osmolyte accumulation in *Medicago truncatula* and *Medicago laciniata* populations. *Comptes Rendus Biologies* 333: 205- 213.
- Zhu JK (2001). Plant salt tolerance. *Trends in Plant Sci.*, 6: 66- 71.